




Contents lists available at ScienceDirect

Middle East Fertility Society Journal

journal homepage: www.sciencedirect.com



Discussions

No cytotoxic effects from application of pentoxifylline to spermatozoa on subsequent pre-implantation embryo development in mice

Mohammad Ali Khalili^a, Azam Agha-Rahimi^{a,*}, Fatemeh Sadeghian^a, Iman Halvaei^b^a Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran^b Department of Anatomical Sciences, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran

ARTICLE INFO

Article history:

Received 21 July 2016

Accepted 4 December 2016

Available online xxx

Keywords:

Pentoxifylline

Fertilization

Blastocyst

DNA fragmentation index

Mouse

ABSTRACT

The aim was to assess the effect of spermatozoa exposed to PTX on the rates of fertilization and embryo development and apoptotic cells within blastocysts in an animal model. Mice Oocytes were inseminated with spermatozoa exposed to 3.6 mmol PTX for 30 min, or with neat spermatozoa. Then fertilization and embryo development rate, blastocyst formation and quality, as well as total cell number of blastocyst, and DNA fragmentation index (DFI) in blastocysts were surveyed in both groups. Fertilization and embryo development rate were similar between the groups. The rates of blastocyst formation did not differ significantly between control and PTX groups (52.4% vs. 51.8%). The average of total cell count in blastocysts and DFI in control and PTX groups were also insignificant (31.08 ± 1.5 vs. 34.14 ± 1.5 and 9.76 ± 5.0 vs. 11.77 ± 5.4). Application of PTX for enhancing sperm motility does not cause a cytotoxic effect on subsequent embryo development and embryo genome integrity.

© 2016 Middle East Fertility Society. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Successful fertilization depends on the quality of spermatozoa, including the motion characteristics. The diminished sperm motility and poor binding capacity to zona pellucida (ZP) are some causes of male infertility. It has been reported that agents such as (PTX) and other Methylxanthines can noticeably improve sperm motility in in vitro condition [1,2]. These chemicals block cyclic nucleotide phosphodiesterase and increase level of intracellular cAMP, then phosphorylation of tyrosine in sperm tail occurs for its activation [3]. Many studies have investigated the effect of PTX on different aspect of spermatozoa, including motion characteristics and viability [4,5]. However, there are only a few studies about the effect of this agent on in vitro embryo development from spermatozoa that were already exposed to PTX. In this regards, Tournaye and co-workers investigated the impact of washed sperms after exposing to 3.6 mmol PTX on mice embryo formation. Their results expressed that PTX exposed spermatozoa had no adverse effects on fertilization and embryo development rates [6]. Also, Rashidi and associates showed that PTX increased fertilization and 2-cell cleavage rates after IVF in mice, when compared with controls [7]. Since then, no other reports have been presented

regarding the efficacy as well as the safety of PTX on generation of mammalian embryos in vitro.

One important physiological phenomenon in reproductive biology is cell apoptosis. During apoptosis, unwanted or useless cells are generally eliminated during development and other normal biological processes. Regulation of apoptosis is very important, especially in pre-implantation embryo development that contains only few vital cells. Inner cell mass (ICM) within blastocysts has been considered as an indicator of embryo quality and loss of cells within ICM below a critical threshold can compromise subsequent fetal development [8,9]. Also, total blastocyst cell numbers is correlated with embryo implantation potential in process of reproduction [10]. Therefore, the aim of this experimental study was to assess the effect of in vitro exposure of spermatozoa to PTX on the rates of fertilization and embryo development as well as apoptotic cells within blastocysts in an animal model.

2. Materials and methods

This study was approved by our institute research committee. Reported experiments were performed according to the appropriate ethical and legal standards.

2.1. Collection and preparation of oocytes

Female Balb/C mice aged 8–10 weeks were induced for superovulation by i.p. injection of 10 IU pregnant mare serum

Peer review under responsibility of Middle East Fertility Society.

* Corresponding author.

E-mail address: 63rahimi@gmail.com (A. Agha-Rahimi).

<http://dx.doi.org/10.1016/j.mefs.2016.12.001>

1110-5690/© 2016 Middle East Fertility Society. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: M.A. Khalili et al., No cytotoxic effects from application of pentoxifylline to spermatozoa on subsequent pre-implantation embryo development in mice, Middle East Fertil Soc J (2016), <http://dx.doi.org/10.1016/j.mefs.2016.12.001>

gonadotropin (PMSG, Sigma, USA). Approximately 48 h later, they were injected with 10 IU of human chorionic gonadotrophin (hCG, Serono, Swiss; i.p.) to trigger ovulation. After 15 h, the oviducts were excised under aseptic conditions and the cumulus-oocyte complexes (COCs) were collected in IVF medium (vitrolife, Sweden). They were cultured in IVF medium (vitrolife, Sweden) in an atmosphere of 5% CO₂ in air at 37 °C. The oocytes were randomly allocated into two groups of control and PTX. In control group, oocytes were inseminated with sperm suspension only. However, in PTX group, insemination was done with sperm suspension exposed to 3.6 mmol PTX.

2.2. Sperm preparation

Male Balb/C mice (10–14 weeks) were killed, and the cauda epididymis were removed immediately. For control group, epididymis was dissected in pre-warm Ham' F10 medium containing 5% serum albumin. For PTX group, it was dissected in aforementioned medium supplemented with 3.6 mmol PTX. Both suspensions were incubated for 30 min to ensure sperm capacitation. Then, for washing, 5 ml Ham' F10 containing 5% serum albumin was added to both suspensions and centrifuged in 300g for 5 min. The supernatants were discarded and the pellets were washed again. The final pellets were used for MII oocytes insemination.

2.3. IVF and embryo cultures

A total of 221 MII oocytes were collected from fallopian tubes of mice, then divided randomly to either the control or PTX groups. For IVF, the oocytes were exposed to spermatozoa for 5 h. Then, oocytes in both groups were washed and cultured in G1 medium (vitrolife, Sweden) for 48 h. The cleaved embryos were cultured in G2 medium (vitrolife, Sweden) until day 5. The rates of fertilization, cleavage, and blastocyst formations were evaluated and compared in both groups. Blastocyst grading was done according to Gardner protocol [11]: (1) early blastocyst: the blastocoels is less than half the embryo volume; (2) blastocyst: the blastocoels is greater than half of the embryo volume; (3) full blastocyst: the blastocoel occupy more than half the volume of the embryo; (4) expanded blastocyst: the blastocoel volume is larger than early embryo and the zona is thinning; (5) hatching blastocyst: the blastocyst has started to exit though the zona; (6) hatched blastocyst: the blastocyst has completely exited the ZP.

2.4. Assessment of apoptosis

For fixation, embryos were incubated in 3.7% paraformaldehyde/PBS for 1 h at room temperature (RT). The embryos were washed in PBS/PVP twice and incubated in 0.5% Triton X-100/PBS for 1 h at RT. After washing in PBS/PVP, the embryos were incubated in 10–15 ml dUTP-FITC labeling mix (Roche, Germany) for 10 min at RT. Then, these were incubated in 10–15 ml of TUNEL mix (Roche, Germany) for 1 h/37 °C in dark. After washing, for nuclear counter stain, embryos were exposed to PBS with Hoechst stain (Sigma). Finally, washing were repeated again and embryos were analyzed immediately in drops of glycerol on glass slides with conventional fluorescence microscopy [11].

2.5. Statistical analysis

SPSS 19 was used for statistical analysis. Differences between data for comparison of the two groups were analyzed by chi-square test and independent *t*-test. P value < 0.05 was statistically significant.

3. Results

3.1. Fertilization and embryo development

The findings generated from this study showed that the rates of fertilization and embryo development were similar between the control and PTX groups. Also, the rates of blastocyst formations did not differ significantly between two groups (Table 1). However, the data demonstrated that both early and expanded blastocyst formations were significantly higher in PTX group when compared with control (Table 2). Fig. 1 shows blastocysts grading in this study.

3.2. Blastocyst cell count and apoptosis

The average of total cell counts in blastocyst stage between two groups were insignificant. However, the DFI was higher in PTX group when compared with control, but this difference was insignificant (Table 3). Fig. 2 shows Hoechst and TUNEL staining of

Table 1

Comparison of fertilization, embryo development and blastocyst rates in control and PTX group.

IVF data	Control	PTX	P value
MI I Oocyte (No.)	110	111	
Fertilization (%)	103 (93.6)	108 (97.2)	0.2
4 Cell embryo (%)	91 (88.3)	92 (85.1)	0.5
Morola (%)	79 (76.6)	82 (75.9)	1.0
Blastocyst (%)	54 (52.42)	56 (51.8)	1.0

Table 2

Comparisons of blastocysts grading in control and PTX groups.

Group	Early	Full	Expand	Hatched
Control (%)	31.4	35.18	11.11	22.2
PTX (%)	7.1	33.9	33.9	25
P value	0.001	1	0.005	0.8




Fig. 1. Stereo photomicrograph of mouse blastocysts in different stages of development.

Download English Version:

<https://daneshyari.com/en/article/8783298>

Download Persian Version:

<https://daneshyari.com/article/8783298>

[Daneshyari.com](https://daneshyari.com)