Accepted Manuscript

Title: Low anti-müllerian hormone concentration is associated with increased risk of embryonic aneuploidy in women of advanced age

Author: Xiao Jiang, Junhao Yan, Yan Sheng, Mei Sun, Linlin Cui, Zi-Jiang Chen

PII: S1472-6483(18)30189-5

DOI: https://doi.org/10.1016/j.rbmo.2018.04.039

Reference: RBMO 1930

To appear in: Reproductive BioMedicine Online

Received date: 1-9-2017 Revised date: 28-3-2018 Accepted date: 4-4-2018



Please cite this article as: Xiao Jiang, Junhao Yan, Yan Sheng, Mei Sun, Linlin Cui, Zi-Jiang Chen, Low anti-müllerian hormone concentration is associated with increased risk of embryonic aneuploidy in women of advanced age, *Reproductive BioMedicine Online* (2018), https://doi.org/10.1016/j.rbmo.2018.04.039.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Short title: Association of AMH with embryonic aneuploidy rate in women of advanced age

Low anti-Müllerian hormone concentration is associated with increased risk of embryonic aneuploidy in women of advanced age

Xiao Jiang a,b, Junhao Yan a,b, Yan Sheng a,b, Mei Sun a,b, Linlin Cui a,b,*, Zi-Jiang Chen a,b,c,* a Centre for Reproductive Medicine, Shandong Provincial Hospital Affiliated to Shandong University; The Key Laboratory of Reproductive Endocrinology (Shandong University), Ministry of Education, 250001 Jinan, China; b National Research Centre for Assisted Reproductive Technology and Reproductive Genetics, 25001 Jinan, China; c Centre for Reproductive Medicine, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University; Shanghai Key Laboratory for Assisted Reproduction and Reproductive Genetics, 200000 Shanghai, China

* Corresponding authors. *E-mail addresses*: chenzijiang@hotmail.com/chen2); fdclear3@126.com (Linlin Cui).

Key message

Low AMH concentration was associated with increased risk of embryo aneuploidy in women of advanced age, but not in young subjects, indicating that embryo aneuploidy risk was mainly due to advanced age and related changes in oocytes and not to ovarian reserve in isolation.



Download English Version:

https://daneshyari.com/en/article/8783717

Download Persian Version:

https://daneshyari.com/article/8783717

<u>Daneshyari.com</u>