




Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Original Article

Obstetric outcomes of twin pregnancies at advanced maternal age: A retrospective study



Caixia Zhu, Malie Wang, Gang Niu, Juan Yang, Zilian Wang*

Department of Obstetrics and Gynecology, The First Affiliated Hospital, Sun Yat-sen University, Guangzhou 510080, China

ARTICLE INFO

Article history:
Accepted 10 August 2017

Keywords:
Obstetric outcomes
Twin pregnancies
Advanced maternal age

ABSTRACT

Objective: To evaluate obstetric outcomes in twin pregnancies of advanced maternal age (≥ 35 years).
Materials and methods: A retrospective study involved 470 twin pregnancies in a single center from Sep. 1, 2012 to Mar. 31, 2015. Clinical characteristics and obstetric outcomes were recorded and compared among twin pregnancies who were classified as follows: age 20–29, 30–34, 35–39 and ≥ 40 years.
Results: The incidence of gestational diabetes (age 20–29 years 15.8%; 30–34 years 24.3%; 35–39 years 30.4%; ≥ 40 years 57.1%; $p = 0.004$) and premature delivery (20–29 years 58.6%; 30–34 years 69.1%; 35–39 years 72.2%; ≥ 40 years 85.7%; $p = 0.001$) significantly increased with increasing age whereas spontaneous abortion (20–29 years 27.6%; 30–34 years 11.6%; 35–39 years 11.4%; ≥ 40 years 0.0%; $p = 0.021$) decreased in twin pregnancies of advanced maternal age. In addition, the rate of postpartum hemorrhage increased almost continuously by age and advanced maternal age was described as a risk factor for postpartum hemorrhage (age 35–39, adjusted OR 3.377; 95% confidence interval 1.729–6.598; $p < 0.001$; age ≥ 40 , adjusted OR 10.520; 95% CI 1.147–96.492; $p = 0.037$). However, there was no significant difference between advanced maternal age and adverse neonatal outcomes.
Conclusion: In twin pregnancies, advanced maternal age experienced significant higher risk of postpartum hemorrhage, gestational diabetes and premature delivery. Neither adverse neonatal outcomes nor stillbirth was significantly associated with maternal age.

© 2018 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The incidence of twin gestation has risen over the past decades in China, owing to the use of assisted reproductive technology (ART), and is associated with high rate of stillbirth and prematurity [1]. Infants born from twin pregnancies consist approximately 20.0% of perinatal and neonatal morbidity and mortality [2]. It is well known that monochorionic-diamniotic (MCDA) twins represent higher rate of neonatal morbidity than dichorionic-diamniotic (DCDA) twins, which is attributed to placental vascular complications, selective intrauterine

growth restriction (sIUGR), maternal obstetric complications and perinatal complications in consequence of prematurity [3]. Considering the low incidence of monochorionic-monoamniotic (MCMA) twins, we only focused on DCDA twins and MCDA twins in this study.

Advanced maternal age, defined as being ≥ 35 years at delivery, is associated with various maternal obstetric complications. Higher incidence of spontaneous abortion, preeclampsia, preterm delivery and stillbirth in singleton pregnancies who aged ≥ 35 years [4]. It is well known that advanced maternal age was identified as an independent risk factor for perinatal complications in singleton pregnancy [5]. To date, much fewer studies focused on twin pregnancies aged ≥ 35 years. Accumulating evidences based on singleton pregnancy. Consequently, the effect of advanced maternal age on twin pregnancies is still unclear.

Recently, there is a consistent trend towards delayed child-bearing in China. Management of twin pregnancies at advanced maternal age remains a challenge in obstetric practice. The purpose of this study was to investigate the associations between maternal age and risk maternal–fetal outcomes in twin pregnancies.

Abbreviations: ART, Assisted reproductive technology; MCDA, Monochorionic-diamniotic; DCDA, Dichorionic-diamniotic; sIUGR, Selective intrauterine growth restriction; PROM, Premature rupture of membrane; GDM, Gestational diabetes mellitus; NICU, Neonatal intensive care unit; RDS, Respiratory distress syndrome; NEC, Necrotizing enterocolitis.

* Corresponding author.

E-mail addresses: cxzhu88@aliyun.com (C. Zhu), 13760649352@126.com (M. Wang), niugdr@163.com (G. Niu), yangjuan1208@126.com (J. Yang), wangzilian2005@aliyun.com (Z. Wang).

<https://doi.org/10.1016/j.tjog.2017.12.010>

1028–4559/© 2018 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Materials and methods

This was a retrospective cohort study of patients with DCDA twins and MCDA twins at a single tertiary center between Sep. 1, 2012 and Mar. 31, 2015.

Patients aged over 20 years with twin pregnancies were included in the study.

Exclusion criteria included MCMA twins, multiple gestation, and twin pregnancies who underwent selective fetal termination at the first trimester. Chorionicity of twin gestation was assigned prenatally by ultrasound and was confirmed by the review of placental pathology. The gestation age was determined by the patients' last menstrual period or date of fertilization in cases of assisted reproduction. We excluded pregnant women aged younger than 20 years old and complained with chronic diseases, including chronic hypertension or collagen vascular disease. This study reduced in accordance with the appropriate clinical and experimental ethical guidelines and was approved by Committee of the First Affiliated Hospital, Sun Yat-sen University. The study was reported with consent from the pregnancies and their relatives.

A total of 470 twin pregnancies, including 330 DCDA twins and 140 MCDA twins, were enrolled in the study. We divided this population into four groups according to the maternal age: 20–29 years, 30–34 years, 35–39 years and ≥ 40 years.

The primary outcome was postpartum hemorrhage. In addition, the secondary obstetric outcome included: Selective intrauterine growth restriction (sIUGR), premature rupture of membrane (PROM), premature delivery, preeclampsia, gestational diabetes mellitus (GDM) and stillbirth. sIUGR was defined as estimated fetal weight (EFW) <10th percentile matched for gender and twin

gestation in the local population and co-twin with EFW >10th percentile. Stillbirth in this study was defined as two fetus died in uterine.

Neonatal outcomes, including neonatal weight, 1-min Apgar score, admission to the neonatal intensive care unit (NICU) rate and incidence of adverse neonatal outcomes, were examined only in the pregnancies who delivered two living newborn. Adverse neonatal outcomes contained respiratory distress syndrome (RDS), jaundice, sepsis, neonatal asphyxia, hypoglycemia, necrotizing enterocolitis (NEC) and so on.

Statistic analysis

All analyses were reported using SPSS version 22.0 (IBM SPSS v 22.0, Armonk, NY). The continuous or dichotomous variables were compared between these groups via the one-way ANOVA. Chi-square test and Fisher's exact test were used to compare the categorical viable. Multivariable logistic regression analysis was performed to determine whether maternal age was significantly independent risk of postpartum hemorrhage and adverse neonatal outcomes. A *p* value of <0.05 was considered significant.

Results

Among 470 twin pregnancies, 203 (43.2%) cases were 20–29 years, 181 (38.5%) cases were 30–34 years, 79 (16.8%) cases were 35–39 years and 7 (1.5%) cases were ≥ 40 years. The maternal characteristic across age groups (Table 1) presented that high level of education was most frequent in group (age 30–34 years; *p* = 0.002). Assisted reproductive technology (ART) was taken more frequently in group (age 35–39, 81.0%; *p* < 0.001). Furthermore, more DCDA twins were showed in group (age 35–39, 79.7%; *p* < 0.001).

As shown in Table 2, the incidence of GDM significantly increased with increasing age (20–29 years 15.8%; 30–34 years 24.3%; 35–39 years 30.4%; ≥ 40 years 57.1%; *p* = 0.004), whereas PROM decreased with increasing age (20–29 years 20.7%; 30–34 years 20.4%; 35–39 years 17.7%; ≥ 40 years 0.0%; *p* = 0.553). In addition, the rate of postpartum hemorrhage increased almost continuously by age: 20–29 years 22.1%; 30–34 years 17.8%; 35–39 years 62.1%; ≥ 40 years 83.3%; *p* < 0.001. There was no significant difference of risk of stillbirth in different age (*p* = 0.133).

Neonatal outcomes according to maternal age at delivery were presented in Table 3. The groups were similar in term of gestational age and birth weight. Likewise, the incidence of adverse neonatal outcome among different maternal age were similar (20–29 years 34.6%; 30–34 years 30.3%; 35–39 years 34.1%; ≥ 40 years 33.3%; *p* = 0.706). Neonatal respiratory distress syndrome (RDS) occurred in 40 women aged 20–29 years, 22 women aged 30–34 years and 8

Table 1
Demographic and obstetrical characteristics (N = 470).

Age (years)	20–29 y	30–34 y	35–39 y	≥ 40 y	p-value
Number	203 (43.2%)	181 (38.5%)	79 (16.8%)	7 (1.5%)	
Education					0.002
Low level	68 (33.5%)	31 (17.1%)	17 (21.5%)	3 (42.9%)	
High level	135 (66.5%)	150 (82.9%)	62 (78.5%)	4 (57.1%)	
Parity					0.212
Nulliparous	154 (75.9%)	132 (72.9%)	51 (64.4%)	6 (85.7%)	
Multiparous	49 (24.1%)	49 (27.1%)	27 (35.4%)	1 (14.3%)	
Conception					
Natural	123 (60.6%)	48 (26.5%)	15 (19.0%)	2 (28.6%)	<0.001
ART*	80 (39.4%)	133 (73.5%)	64 (81.0%)	5 (71.4%)	
Previous cesarean	16 (7.9%)	14 (7.7%)	6 (16.7%)	0 (0.0%)	0.897
Chorionicity					
MCDA#	87 (42.9%)	38 (21.0%)	16 (20.3%)	2 (28.6%)	<0.001
DCDA**	116 (57.1%)	143 (79.0%)	63 (79.7%)	5 (71.4%)	

*ART: assisted reproductive technology; #MCDA: monochorionic-diamniotic twins; **DCDA: dichorionic-diamniotic twins.

Table 2
Obstetric and clinical outcomes (N = 470).

Age (years)	20–29 y	30–34 y	35–39 y	≥ 40 y	p-value
Preeclampsia	14 (6.9%)	12 (6.6%)	7 (8.9%)	2 (5.7%)	0.342
GDM*	32 (15.8%)	44 (24.3%)	24 (30.4%)	4 (57.1%)	0.004
sIUGR***	32 (15.8%)	20 (11.0%)	9 (11.4%)	0 (0.0%)	0.360
PROM#	42 (20.7%)	37 (20.4%)	14 (17.7%)	0 (0.0%)	0.553
Stillbirth	14 (6.9%)	6 (3.3%)	1 (1.3%)	0 (0.0%)	0.133
Congenital malformations	24 (11.8%)	11 (6.1%)	13 (16.5%)	1 (14.3%)	0.064
Postpartum hemorrhage	30 (22.1%)	28 (17.8%)	41 (62.1%)	5 (83.3%)	<0.001
Gestational age at delivery					
<28 wk	56 (27.6%)	21 (11.6%)	9 (11.4%)	0 (0.0%)	0.021
28–37 wk	119 (58.6%)	125 (69.1%)	57 (72.2%)	6 (85.7%)	0.001
≥ 37 wk	28 (13.8%)	35 (19.3%)	13 (16.5%)	1 (14.3%)	0.324

*GDM: gestational diabetes; ***sIUGR: selective intrauterine growth restriction; #PROM: premature rupture of membrane.

Forbidden

You don't have permission to access this resource.

Apache/2.4.25 (Debian) Server at daneshyari.com Port 443