



Original Article

Preliminary report of altered insulin secretion pattern in monochorionic twin pregnancies complicated with selective intrauterine growth restriction



Yao-Lung Chang^{a, b, *}, Tzu-Hao Wang^{a, b}, Seham M. Abufraijeh^c, Shuenn-Dyh Chang^{a, b}, An-Shine Chao^{a, b}, Peter C.C. Hsieh^{a, b}

^a Department of Obstetrics and Gynecology, Chang Gung Memorial Hospital, Linkou, Taoyuan, Taiwan

^b Department of Obstetrics and Gynecology, Chang Gung University College of Medicine, Tao-Yuan, Taiwan

^c Department of Obstetrics and Gynecology, College of Medicine, Mutah University Al-Karak, Jordan

ARTICLE INFO

Article history:

Accepted 10 November 2015

Keywords:

growth restriction
insulin
monozygotic twins

ABSTRACT

Objective: Fetuses with intrauterine growth restriction (IUGR) have adaptive hormonal changes including changes in insulin, which may increase their future risks for developing diabetes mellitus. This study compared cord blood insulin concentrations in IUGR and appropriate for gestational age (AGA) fetuses in a monozygotic (MC) twin model.

Materials and methods: Ten pairs were classified as selective IUGR (sIUGR) based on having one twin weight below the 10th percentile and with an intertwin birth weight discordance > 20%. Fourteen pairs without IUGR were included as a comparison group. Pregnancies with twin–twin transfusion syndrome, congenital structural malformations, and genetic abnormalities were excluded. Insulin and glucose concentrations were measured in cord venous blood at the time of delivery.

Results: Cord blood insulin concentrations of sIUGR fetuses were significantly lower than those of AGA counterpart fetuses in MC twins affected by sIUGR (5.1 ± 4.1 mU/L, range: 0.7–9.9 mU/L for sIUGR fetuses and 12.2 ± 7.6 mU/L, range: 3.5–23.7 mU/L for AGA fetuses, $p = 0.019$). No significant difference in insulin concentrations between larger and smaller fetuses in MC twins without IUGR was observed. Insulin concentration was inversely correlated with gestational age of delivery in all fetuses except in those with sIUGR. We did not find any difference in cord blood glucose concentrations between the two fetuses in both groups.

Conclusion: Our data show reduced insulin secretion and loss of the physiological decline in concentration over time as gestational age increases in fetuses with sIUGR compared to AGA counterparts.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Fetal insulin can be detected as early as Week 8 of gestation. It is produced by the fetal pancreas since no maternal insulin can cross the placenta to the fetal circulation due to the large molecular weight of insulin [1].

Previous studies that evaluated fetal insulin concentrations in growth-restricted fetuses have shown mixed results. Some studies

showed reduced fetal insulin concentrations in growth restricted fetuses compared to appropriate for gestational age (AGA) [2–4] fetuses, while others revealed no change [5] in either singleton [2–4,6] or twin pregnancies [5].

Many studies have suggested a relationship between fetal low birth weight and increased risk of impaired glucose tolerance later in life. Adults with a history of intrauterine growth restriction (IUGR) have higher insulin levels at baseline and post 75-g oral glucose tolerance test (OGTT) [7–9], indicating peripheral insulin resistance. The hypothesis of “thrifty phenotype” proposes that a fetus suffering from IUGR would adapt to a poorer intrauterine environment by optimizing the use of a reduced nutrient supply to ensure survival [10]. These changes might aid survival of the fetus,

* Corresponding author. Department of Obstetrics and Gynecology, Chang Gung Memorial Hospital, Linkou Medical Center, 5, Fu-Shin Street, Kweishan, Taoyuan, 333, Taiwan.

E-mail address: j12054@cgmh.org.tw (Y.-L. Chang).

but they become a liability in situations of nutritional abundance, and would thus have a higher chance to cause metabolic diseases such as type 2 diabetes mellitus [11].

Selective IUGR (sIUGR), defined as one IUGR twin with a co-twin who is AGA, occurs in ~12% of twin pregnancies [12,13]. MC twin pregnancy can be a good model for studying the effects of IUGR, since those fetuses have an identical genetic makeup and share the same maternal surrounding. That would leave the placental factor or placental perfusion as a sole useful variable for evaluation of the effects of IUGR on investigated targets.

This study was designed to evaluate the insulin concentration in monochorionic (MC) twins with sIUGR, with their AGA twin counterparts as the internal control group, and a separate MC twin pregnancy without IUGR as an external control group.

Materials and methods

The study participants were MC twins delivered between February 2013 and December 2014, in Chang Gung Memorial Hospital, Linkou Branch; a tertiary referral hospital in Taiwan. Pregnancies with twin–twin transfusion syndrome (TTTS), and congenital structural malformations and genetic abnormalities of fetuses were excluded. The definition of TTTS is as for the Quintero' Staging System [14]. The pregnancies included were then divided into two groups: Group 1 was composed of pregnancies with MC twins complicated by sIUGR, which is defined as an estimated fetal weight for the smaller twin below the 10th percentile under a standard singleton pregnancy birth weight chart and a birth weight discordance > 20% [15] between the sIUGR twins and their AGA twin counterparts; and Group 2 included pregnancies with MC twins without IUGR.

Birth weight discordance was calculated as the difference between the fetal weight of the larger twin and the smaller twin divided by the fetal weight of the larger twin:

$$[\text{body weight of larger twin (AGA)} - \text{body weight of smaller twin (sIUGR)}] / \text{body weight of larger (AGA) twin} \times 100\%.$$

First trimester or early second trimester ultrasound done by a trained ultrasonographer was used to determine chorionicity, and was confirmed by examination of the placentae after delivery.

Table 1
Characteristics of MC twins with and without sIUGR.

	MC twins with sIUGR (n = 10)	MC twins without IUGR (n = 14)	p
Maternal age (y)	31.6 ± 3.3 (27.3–38.4)	32.9 ± 2.4 (30.2–38.7)	0.279 ^a
Gravidity (range)	2 (1–3)	2 (1–3)	0.666 ^b
Parity (range)	1 (1–2)	1 (1–3)	0.841 ^b
GA at delivery (wk)	32.6 ± 2.0 (28.7–35.3)	35.8 ± 1.8 (31.3–38.1)	0.001 ^a
Birth weight of AGA or larger twin (g)	1791 ± 449 (1000–2500)	2331 ± 396 (1470–2880)	< 0.001 ^a
Birth weight of sIUGR or smaller twin (g)	1197 ± 481 (476–1840)	2175 ± 408 (1230–2610)	0.008 ^a
Birth weight discordance (%)	34.5 ± 13.6 (20.1–52.4)	6.7 ± 5.4 (0.8–18.9)	< 0.001 ^a

Data are expressed as mean ± standard deviation (range) or median (range).

AGA = appropriate for GA; GA = gestational age; IUGR = intrauterine growth restriction; MC = monochorionic; sIUGR = selective intrauterine growth restriction.

^a Two-sample Student's *t* test.

^b Mann–Whitney *U* test.

Table 2
Insulin concentrations between the two fetuses of MC twins.

	Insulin concentration of IUGR twin (smaller) (mU/L)	Insulin concentration of AGA twin (larger) (mU/L)	p
MC twins with sIUGR (n = 10)	5.1 ± 4.1 (0.7–9.9)	12.2 ± 7.6 (3.5–23.7)	0.019 ^a
MC twins without IUGR (n = 14)	5.6 ± 5.2 (0.6–19.6)	6.3 ± 4.7 (2.2–20.2)	0.32 ^a

Data are expressed as mean ± standard deviation (range).

The insulin concentration was measured by ARCHITECT Insulin Reagent Kit using chemiluminescence immunoassay provided by Abbott (Wiesbaden, Germany).

AGA = appropriate for gestational age; IUGR = intrauterine growth restriction; MC = monochorionic; sIUGR = selective intrauterine growth restriction.

^a Two-sample Student's *t* test.

Umbilical cord venous blood samples were collected immediately after delivery for both groups, followed by measuring insulin and glucose concentrations using chemiluminescence immunoassay and enzymatic methods, respectively. Both tests were performed by the hospital central laboratory for commercial practice. The tests were provided upon request.

Informed consent was obtained from parents, and the study protocol was approved by the Institutional Review Board (IRB 101-4803A3).

Statistical analysis was conducted with SPSS for Windows version 11.0 (SPSS Inc., Chicago, IL, USA). Two-sample Student's *t* test or Mann–Whitney *U* test was used to compare between groups for the continuous variables. Data within twin pair was compared with the paired-sample *t* test. Correlation of serum insulin with gestational age (GA) was calculated by Pearson correlation test with a *p* value < 0.05 regarded as statistically significant.

Results

Among 24 MC twin pregnancies delivered between February 2013 and December 2014, 10 pregnancies were complicated by sIUGR and 14 were without IUGR. Since our hospital is a tertiary referral center, the percentage of pregnancies with sIUGR was high. The demographics and baseline characteristics of twins are shown in Table 1. Twins with sIUGR were delivered earlier than twins without IUGR. Mean GA at delivery was 32.6 ± 2.0 weeks (range: 28.7–35.3 weeks) in twins with sIUGR compared to 35.8 ± 1.8 weeks (range: 31.3–38.1 weeks) for AGA twins (*p* = 0.001). In the 10 cases with sIUGR, two suffered from fetal distress with repeated fetal heart beat deceleration and they thus were delivered at GA of 28 weeks 5 days and 31 weeks 2 days, respectively. Three cases were scheduled for delivery at 34 weeks after consultation with patients and their family about the risk of intrauterine demise of the sIUGR twin, and the subsequent detrimental effect that might happen to the AGA twins following their death. Pairs with sIUGR demonstrated a significantly higher discordance in birth weight compared with AGA pairs: 34.5 ± 13.6% versus 6.7 ± 5.4%, respectively (*p* < 0.001).

Download English Version:

<https://daneshyari.com/en/article/8784654>

Download Persian Version:

<https://daneshyari.com/article/8784654>

[Daneshyari.com](https://daneshyari.com)