



Original Article

Incidence and lifetime risk of uterine corpus cancer in Taiwanese women from 1991 to 2010



Jerry Cheng-Yen Lai^a, Chia-Sui Weng^b, Sheng-Miauh Huang^c, Nicole Huang^{d,e},
Yiing-Jenq Chou^a, Chien-Chen Wang^f, Kung-Liahng Wang^{b,g,h,*}

^a Institute of Public Health & Department of Public Health, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^b Department of Obstetrics and Gynecology, Mackay Memorial Hospital and Mackay Medical College, Taipei, Taiwan

^c Department of Nursing, Mackay Medical College, New Taipei City, Taiwan

^d Institute of Hospital and Health Care Administration, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^e Department of Education and Research, Taipei City Hospital, Taipei, Taiwan

^f Department of Medicine, Medical University of Lublin, Lublin, Poland

^g Department of Nursing, Mackay Junior College of Medicine, Nursing, and Management, Taipei, Taiwan

^h Department of Obstetrics and Gynecology, Mackay Memorial Hospital Taitung Branch, Taitung, Taiwan

ARTICLE INFO

Article history:

Accepted 18 September 2015

Keywords:

endometrial cancer
incidence
lifetime risk
population surveillance
Taiwan

ABSTRACT

Objective: Although uterine corpus cancer has been the most common malignancy of the female genital tract in many countries, the lifetime risk of this cancer has not yet been determined among Taiwanese women. The purpose of the study was to describe the change in incidence and the lifetime risk of uterine corpus cancer over a 20-year period from 1991 to 2010 in Taiwan.

Materials and methods: We conducted a population-based registry study using the released database (available online) from the Taiwan Cancer Registry.

Results: A total of 15,542 women newly diagnosed with uterine corpus cancer were included in this study. The total number of this cancer increased by 5.7-fold from 1991 to 2010. The annual age-specific rate nearly doubled during the past decade (2001–2010) when compared with the previous decade (1991–2000). Incidence rates were highest in women aged 50–59 years, and increasing incidence rates were observed in each age strata starting from 40 years to 85 years and more, after the year 2000. The lifetime risk of being diagnosed with uterine corpus cancer was 0.39% in 1991–1995, 0.54% in 1996–2000, 0.73% in 2001–2005, and 1.12% in 2006–2010 among Taiwanese women.

Conclusion: According to the observed changes in incidence rate, the burden of uterine corpus cancer in the general female population is expected to increase in the near future. From a public-health perspective, care providers should develop strategies for the prevention, early detection, and intervention to reduce the rapidly increasing incidence of uterine corpus cancer in Taiwan.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Uterine corpus cancer occurs predominately in postmenopausal women aged 50–60 years [1]. Endometrial carcinoma is the most common histologic type, which accounts for approximately 80% of the corpus malignancy [2]. According to the online GLOBOCAN 2012 database [3], the age-standardized incidence rates (ASR) of uterine corpus cancer are as high as 16.3–19.5 per 100,000 person-years

in several Western countries (USA, Canada, and Norway), whereas a low rate of 2–11.2 per 100,000 person-years are seen in many Asian countries (China, Korea, India, Thailand, Malaysia). In the USA, the Surveillance, Epidemiology, and End Results data of nine registries indicate that the incidence of uterine corpus cancer remains relatively stable over the past 30 years (ASR of 20.0 in 1980 vs. 18.3 per 100,000 women/year in 2007) [4]. In contrast, uterine corpus cancer is the most rapidly increasing gynecologic malignancy among Taiwanese women. During the same timeframe, the ASR of uterine corpus cancer has nearly septupled from 1.69 in 1980 to 11.36 per 100,000 women/year in 2010 [5]. In 2010, uterine corpus cancer ranked as the sixth most common cancer in women, and accounted

* Corresponding author. Mackay Memorial Hospital, Taitung Branch, Number 1, Lane 303, Changsha Street, Taitung City 950, Taiwan.

E-mail address: KL421229@ms6.hinet.net (K.-L. Wang).

for roughly 4.4% of all newly diagnosed female malignancies, with 1752 new cases.

The lifetime risk of developing cancer is commonly used to provide a risk estimate in a population, which renders the proportion of the population expected to develop the disease over the course of a lifetime [6]. Although uterine corpus cancer receives considerable attention in many developed countries, the lifetime risk of this cancer as a whole has not yet been determined among Taiwanese women. The purpose of this study was to describe the change in incidence and the lifetime risk of uterine corpus cancer over a 20-year period from 1991 to 2010 in Taiwan.

Materials and methods

Data on uterine corpus cancer incidence, age-specific, and age-standardized (to the world standard 2000 population) incidence rates per 100,000 women/year were obtained for the years 1991–2010 from the released database (available online) at the official website of Taiwan Cancer Registry (TCR) [5]. All women were diagnosed with uterine corpus cancer (International Classification of Diseases for Oncology, Field Trial 182 and International Classification of Diseases for Oncology-3 C54). The primary site codes were classified according to the International Classification of Diseases for Oncology, Field Trial edition, and shifted to the third edition version (International Classification of Diseases for Oncology-3) in 2002. TCR is a population-based national cancer registry, established by the Ministry of Health and Welfare in 1979 to estimate cancer incidence among all civilian residents in Taiwan. Hospitals with 50 or more beds throughout the country are required to report diagnostic and surgical information on newly diagnosed cases of cancer to the TCR on an annual basis. Since all data used in this study were online, publicly accessible aggregate datasets that were openly available to anyone, Institutional Review Board approval was not required for this study.

All rates were expressed as per 100,000 women/year (over 1 year of observation) or per 100,000 person-years (over several years of observation). We compared the age-specific rate, estimated annual percent change (EAPC), and the lifetime risk of four calendar periods (1991–1995, 1996–2000, 2001–2005, and 2006–2010). Since the age-specific rate was not available for the period 2006–2010, it was calculated using stratified procedure counts (by 5-year age groups) and corresponding population denominators (from the National Statistics, Republic of China) [7]. In 2010, 49.8% of the Taiwan population (or 11.5 million civilian residents) were women. The EAPC was used to describe the annual percentage change of age-specific rate for each calendar period. By fitting a regression line to the natural logarithm of age-specific rate using calendar year as a regression variable, EAPC was calculated using the exponential formula $[EAPC = 100 * (\exp^m - 1)]$, where “m” was the slope parameter of the regression line and “exp” was the exponential function. Although the cumulative risk calculation proposed by Day (1987) [8] was not directly equivalent to the lifetime risk of developing cancer over a lifetime [6], by setting the truncated upper age band to the average life expectancy in Taiwanese women, the calculated cumulative risk was approximately equivalent to the lifetime risk. Since the average life expectancy among Taiwanese women increased from 76.76 years in 1991 to 82.47 years in 2010 [9], an average of 79 years was used as the average life expectancy of the female population in Taiwan between 1991 and 2010 for the cumulative risk calculation. The cumulative rate expressed as a percentage was first calculated from the cross-sectional age-specific rate until age 79 years. The cumulative risk was then calculated from the cumulative rate using the exponential formula $[\text{cumulative risk} = 1 - \exp^{-\text{cumulative rate}}]$ as proposed by Day (1987) [8]. All calculations were performed using

SAS software version 9.2 (SAS Institute Inc., Cary, NC, USA). All reported *p*-values were based on two-sided tests at the 0.05 level of statistical significance.

Results

A total of 15,542 women were newly diagnosed with uterine corpus cancer in Taiwan between 1991 and 2010 (Table 1). The numbers of uterine corpus cancer increased by 5.7-fold from 262 cases in 1991 to 1752 cases in 2010 (1.4-fold from 1991 to 2000; 1.5-fold from 2001 to 2010). The EAPC from 2001 to 2010 was 14.25 [95% confidence interval (CI): 4.91–24.43], which was almost twice the estimated annual rate between 1991 and 2000. Although the annual increase in rates for the 1991–1995 period (5.01%/year) was not statistically significant (95% CI: 1.56–8.58), there was a significant increase in the EAPC for the other three calendar periods [9.62% (95% CI: 5.54–13.86) in 1996–2000; 8.67% (95% CI: 6.97–10.39) in 2001–2005; and 10.23% (95% CI: 7.13–13.42) in 2006–2010].

The age-standardized incidence rate of uterine corpus cancer increased by 2.8 times from 3.02 cases in 1991 to 11.36 cases per 100,000 women in 2010 (Figure 1). Figure 2 demonstrates a shift in the age distribution of uterine corpus cancer risk between the periods 1991–1995 and 2001–2005. The risk increased with age and peaked at women aged 50–59 years for all periods. However, the next highest peak was at women aged 75–79 years during the period 1991–1995, but shifted to those aged 80–84 year after year 2000. In other words, more elderly women were diagnosed of uterine corpus cancers in the periods 2001–2005 and 2006–2010 than during the periods 1991–1995 and 1996–2000. In 2010, approximately 65% of all incident cases were diagnosed among women aged 40–59 years (41.4% among women aged 50–59 years and 23.5% among those aged 40–49 years), followed by 26.7% among women aged 60 years or above, and the remaining 8.4% in women aged less than 40 years.

Figure 3 shows the lifetime risk (cumulative risk before age 79 years) of incident cases of uterine corpus cancer in the 1991–1995, 1996–2000, 2001–2005, and 2006–2010 periods, based on the cross-sectional age-specific incidence rates in Table 1. The lifetime risk of being diagnosed with uterine corpus cancer was 0.39%, 0.54%, 0.73%, and 1.12% for 1991–1995, 1996–2000, 2001–2005, and 2006–2010 periods, respectively. In 2010, the lifetime risk of uterine corpus cancer was 1.25% for a 79-year-old woman and 1.35% for women aged ≥ 85 years (Figure 4).

Discussion

This study investigated the changes in incidence and the lifetime risk of uterine corpus cancer in the general female population in Taiwan from 1991 to 2010. Our findings suggested that the incidence of uterine corpus cancer increased by 5.7-fold during the study period. The annual age-specific rate nearly doubled during the past decade (2001–2010) when compared with the previous decade (1991–2000), particularly in the latest period from 2006 to 2010. Incidence rates were highest in women aged 50–59 years, and increasing incidence rates were observed in each age strata starting from 40 years to 85 years and more, after the year 2000. The lifetime risk of being diagnosed with uterine corpus cancer was 0.39% in 1991–1995, 0.54% in 1996–2000, 0.73% in 2001–2005, and 1.12% in 2006–2010 among Taiwanese women.

The majority of endometrial cancers (70–80%) are estrogen-related malignancies [10]. The rapid increase in the incidence and the lifetime risk of uterine corpus cancer are most likely attributed to the increasingly prevalence of both exogenous and endogenous estrogen sources among Taiwanese women over the past 20 years

Download English Version:

<https://daneshyari.com/en/article/8784659>

Download Persian Version:

<https://daneshyari.com/article/8784659>

[Daneshyari.com](https://daneshyari.com)