


Accepted Manuscript



Title: A component of lobular carcinoma in clinically lymph node-negative patients predicts for an increased likelihood of upstaging to pathologic stage III breast cancer

Author: Renae D. Van Wyhe, Abigail S. Caudle, Simona F. Shaitelman, George H. Perkins, Thomas A. Buchholz, Karen E. Hoffman, Eric A. Strom, Benjamin D. Smith, Welela Tereffe, Wendy A. Woodward, Michael C. Stauder

PII: S2452-1094(18)30032-0
DOI: <https://doi.org/10.1016/j.adro.2018.02.007>
Reference: ADRO 175

To appear in: *Advances in Radiation Oncology*

Received date: 23-10-2017
Revised date: 7-2-2018
Accepted date: 14-2-2018

Please cite this article as: Renae D. Van Wyhe, Abigail S. Caudle, Simona F. Shaitelman, George H. Perkins, Thomas A. Buchholz, Karen E. Hoffman, Eric A. Strom, Benjamin D. Smith, Welela Tereffe, Wendy A. Woodward, Michael C. Stauder, A component of lobular carcinoma in clinically lymph node-negative patients predicts for an increased likelihood of upstaging to pathologic stage III breast cancer, *Advances in Radiation Oncology* (2018), <https://doi.org/10.1016/j.adro.2018.02.007>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: A Component of Lobular Carcinoma in Clinically Lymph Node-Negative Patients Predicts for an Increased Likelihood of Upstaging to Pathologic Stage III Breast Cancer

Short title: Invasive Lobular Carcinoma and Risk of Surgical Upstaging

Authors: Renae D. Van Wyhe, MBA^{1,3}, Abigail S. Caudle, MD², Simona F. Shaitelman, MD¹, George H. Perkins, MD¹, Thomas A. Buchholz, MD¹, Karen E. Hoffman, MD¹, Eric A. Strom, MD¹, Benjamin D. Smith, MD¹, Welela Tereffe, MD¹, Wendy A. Woodward, MD, PhD¹, and Michael C. Stauder, MD¹

Author affiliations: Departments of ¹Radiation Oncology and ²Breast Surgical Oncology, The University of Texas MD Anderson Cancer Center, ³Baylor College of Medicine

Corresponding Author:

Michael C. Stauder, MD

Assistant Professor, Radiation Oncology

The University of Texas MD Anderson Cancer Center

1515 Holcombe Blvd., Unit 1202

Houston, TX 77030

mstauder@mdanderson.org

Telephone: 713-563-3764

Fax: 713-563-6940

Conflict of Interest Statement: The authors whose names are presented in this manuscript have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

Acknowledgments: None to include.

Summary: Clinical evaluation is often less precise in determining the extent of disease in invasive lobular carcinoma compared to non-lobular histologies. In clinically node-negative patients with lobular histology, we report a 30% incidence of upstaging to pathologic stage IIIA or higher.

Abstract

Purpose: Physical exam and diagnostic imaging are often less precise in determining the extent of disease in invasive lobular carcinoma (ILC) relative to non-lobular histologies. Anecdotally,

Download English Version:

<https://daneshyari.com/en/article/8784796>

Download Persian Version:

<https://daneshyari.com/article/8784796>

[Daneshyari.com](https://daneshyari.com)