

# Accepted Manuscript

Pre-treatment Neutrophil-to-Lymphocyte Ratio as a predictor for occult cervical metastasis in early Stage (T1-T2 cN0) Squamous Cell Carcinoma of the Oral Tongue



Vincenzo Abbate, Giovanni Dell'Aversana Orabona, Giovanni Salzano, Paola Bonavolontà, Fabio Maglitto, Antonio Romano, Filippo Tarabbia, Mario Turri Zanoni, Federica Attanasi, Alessandro Espedito Di Lauro, Giorgio Iaconetta, Luigi Califano

PII: S0960-7404(18)30014-8

DOI: 10.1016/j.suronc.2018.06.002

Reference: SO 1104

To appear in: *Surgical Oncology*

Received Date: 14 January 2018

Accepted Date: 02 June 2018

Please cite this article as: Vincenzo Abbate, Giovanni Dell'Aversana Orabona, Giovanni Salzano, Paola Bonavolontà, Fabio Maglitto, Antonio Romano, Filippo Tarabbia, Mario Turri Zanoni, Federica Attanasi, Alessandro Espedito Di Lauro, Giorgio Iaconetta, Luigi Califano, Pre-treatment Neutrophil-to-Lymphocyte Ratio as a predictor for occult cervical metastasis in early Stage (T1-T2 cN0) Squamous Cell Carcinoma of the Oral Tongue, *Surgical Oncology* (2018), doi: 10.1016/j.suronc.2018.06.002

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Pre-treatment Neutrophil-to-Lymphocyte Ratio as a predictor for occult cervical metastasis in early Stage (T1-T2 cN0) Squamous Cell Carcinoma of the Oral Tongue**

Vincenzo Abbate<sup>a</sup> MD, Giovanni Dell'Aversana Orabona<sup>a</sup> MD, Giovanni Salzano<sup>a</sup> MD, Paola Bonavolontà<sup>a</sup> MD, Fabio Maglitto<sup>a</sup> MD, Antonio Romano<sup>a</sup> MD, Filippo Tarabbia<sup>a</sup> MD, Mario Turri Zanoni<sup>b</sup> MD, Federica Attanasi<sup>c</sup>, Alessandro Espedito Di Lauro MD<sup>d</sup>, Giorgio Iaconetta<sup>e</sup> MD, Luigi Califano<sup>a</sup> MD.

<sup>a</sup> Maxillofacial Surgery Unit, Department of Neurosciences, Reproductive and Odontostomatological Sciences, University Federico II, Via Pansini 5, 80100 Naples, Italy.

<sup>b</sup> Division of Otorhinolaryngology, Department of Biotechnology and Life Sciences, University of Insubria, Ospedale di Circolo e Fondazione Macchi, Varese, Italy

<sup>c</sup> Department of Statistical Sciences, University La Sapienza, Piazzale Aldo Moro 5, 00185 Rome, Italy.

<sup>e</sup> Division of Dentistry, Department of Neurosciences, Reproductive and Odontostomatological Sciences, University Federico II, Via Pansini 5, 80100 Naples, Italy.

<sup>d</sup> Neurosurgery Unit Department of Medicine, Surgery and Odontoiatrics, University of Salerno, Via Giovanni Paolo II 132, 84084 Fisciano, Salerno, Italy.

Corresponding author: Vincenzo Abbate MD tel: +39 3357266177, fax: +39 0817462179; e-mail address: vincenzo.abbate@unina.it. Institutional Address: Via Sergio Pansini 5, Naples, Italy. Postal code 80131

Brief Title: NLR to stratify patients based on risks of metastatic node

**Key words:** Oral tongue squamous cell carcinoma; NLR; neutrophil to lymphocyte ratio; Pre-treatment hematological marker; Occult metastasis.

### **Conflict of Interest and Source of Funding**

The authors do not declare conflict of interest and did not receive any financial support for the study presented in the manuscript.

Download English Version:

<https://daneshyari.com/en/article/8789679>

Download Persian Version:

<https://daneshyari.com/article/8789679>

[Daneshyari.com](https://daneshyari.com)