Accepted Manuscript



Myeloid cells contribute indirectly to VEGF expression upon hypoxia via activation of Müller cells

Christina Nürnberg, Norbert Kociok, Claudia Brockmann, Timo Lischke, Sergio Crespo-Garcia, Nadine Reichhart, Susanne Wolf, Ria Baumgrass, Sabine A. Eming, Sandra Beer-Hammer, Antonia M. Joussen

PII: S0014-4835(17)30458-X

DOI: 10.1016/j.exer.2017.10.011

Reference: YEXER 7222

To appear in: *Experimental Eye Research*

Received Date: 20 July 2017

Revised Date: 6 September 2017

Accepted Date: 8 October 2017

Please cite this article as: Nürnberg, C., Kociok, N., Brockmann, C., Lischke, T., Crespo-Garcia, S., Reichhart, N., Wolf, S., Baumgrass, R., Eming, S.A., Beer-Hammer, S., Joussen, A.M., Myeloid cells contribute indirectly to VEGF expression upon hypoxia via activation of Müller cells, *Experimental Eye Research* (2017), doi: 10.1016/j.exer.2017.10.011.

Myeloid cells contribute indirectly to VEGF expression upon hypoxia via activation of Müller cells

Christina Nürnberg[1], Norbert Kociok[1], Claudia Brockmann[1], Timo Lischke[2], Sergio Crespo-Garcia[1], Nadine Reichhart[1], Susanne Wolf[3], Ria Baumgrass[2], Sabine A. Eming[4], Sandra Beer-Hammer[5], and Antonia M. Joussen[1]

[1]Department of Ophthalmology, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany

[2]German Rheumatism Research Center Berlin, a Leibniz Institute, Berlin, Germany

[3]Department of Cellular Neuroscience, Max Delbrück Center in the Helmholtz Society, Berlin, Germany

[4]Department of Dermatology, University of Cologne, Germany

[5]Department of Pharmacology and Experimental Therapy and Interfaculty Center of Pharmacogenomics and Drug Research, University of Tübingen, Germany