


Contents lists available at [ScienceDirect](#)

## Acta Orthopaedica et Traumatologica Turcica

journal homepage: <https://www.elsevier.com/locate/aott>

## Return to sport activity after anterior cruciate ligament reconstruction in skeletally immature athletes with manual drilling original all inside reconstruction at 8 years follow-up

Giacomo Placella <sup>a,\*</sup>, Matteo Bartoli <sup>a</sup>, Marco Peruzzi <sup>a</sup>, Andrea Speziali <sup>a</sup>, Valerio Pace <sup>b</sup>, Giuliano Cerulli <sup>a</sup>

<sup>a</sup> Institute of Orthopedics and Traumatology, Catholic University of the Sacred Heart, 'Agostino Gemelli' Hospital, Largo Agostino Gemelli 8, 00168 Rome, Italy

<sup>b</sup> Royal National Orthopaedi Hospital, Stanmore, London, UK

## ARTICLE INFO

## Article history:

Received 19 October 2015

Received in revised form

12 January 2016

Accepted 1 March 2016

Available online xxx

## Keywords:

All-inside

ACL

Athlete

Sports

Adolescent

## ABSTRACT

**Objective:** Anterior cruciate ligament (ACL) tears are common injuries in adolescent athletes, especially in those who bear high stress on their knees due to shearing forces. The goal of the surgical procedures in skeletally immature patients is to restore joint stability avoiding the adverse effects on the growth process.

The aim of this study was to verify the return of the skeletally immature professional athletes to sports in the long-term, following ACL reconstruction with the original all-inside technique and with manual drilling.

**Methods:** This study included 24 athletes (14 boys, 10 girls; mean age: 13.15 years, range: 9–14 years) who had radiographic evidence of open physes, were less than 14 years of age at the time of surgery and those with a minimum follow-up of eight years. All patients completed a questionnaire, the IKDC subjective knee evaluation form, and Tegner Activity Scale. Biomechanical outcomes of the KT-1000 arthrometer, gait analysis, and stabilometric and isokinetic results were also evaluated. A plain radiograph of both lower limbs was taken to obtain a precise measurement of the limb length and mechanical axis angles.

**Results:** The patients returned to sport activities in a mean time of 6.43 months. No rerupture or resurgery due to growth abnormalities was observed. The mean difference in length between the operated and contralateral legs was 0.4 (range: –0.2 to 0.7) cm. The mean side-to-side difference measured with the KT-1000 arthrometer was 5.2 (range: 3.5 to 7) mm in the preoperative and 0.8 (range: 0 to 2.5) mm in the postoperative measurements.

**Conclusion:** In conclusion, the original all-inside technique with manual drilling with a half tunnel and short graft seems to be a very effective technique for the surgical management of ACL injuries in pediatric/adolescent athletes. A good rate of return to sports at pre-injury levels or higher, high patient satisfaction and a decent motor and proprioceptive function is possible as shown by our analysis.

**Level of evidence:** Level IV, Therapeutic study.

© 2016 Turkish Association of Orthopaedics and Traumatology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Anterior cruciate ligament (ACL) tears are common injuries in adolescent athletes, especially in those who bear high stress on their knees due to shearing forces.<sup>1–3</sup> The goal of the surgical procedures in skeletally immature patients is to restore

joint stability avoiding the adverse effects on the growth process.<sup>4</sup>

In order to avoid the risks that may occur with an unstable knee, some authors believe that a change in the young athletes' activity program before starting the standard program could get them ready for skeletal maturity. However, it is clear that prohibiting a young athlete from doing sports is not a feasible solution. Restoring the joint stability is necessary to prevent secondary meniscal and

\* Corresponding author. Let People Move, via G.B. Pontani, 9, Italy. Fax: +39 075 5292165.

E-mail address: [giacomo.placella@gmail.com](mailto:giacomo.placella@gmail.com) (G. Placella).

<http://dx.doi.org/10.1016/j.aott.2016.03.006>

1017-995X/© 2016 Turkish Association of Orthopaedics and Traumatology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

cartilaginous lesions, and allow young athletes to continue with their careers.<sup>5–7</sup>

Clinical studies have shown good functional outcomes and high rates of return to sports after ACL reconstruction among young patients, through different technical possibilities to create bone tunnels<sup>8–11</sup>; the transphyseal technique,<sup>6</sup> the mixed technique,<sup>12</sup> the extraepiphyseal technique<sup>13</sup> and physeal sparing.<sup>14</sup> Most of the authors make their decisions based on the presence or absence of the patient's residual growth plate.<sup>4</sup>

Mechanical and biological reasons may affect the bone growth. The mechanical reasons might arise from the mechanism of fixation and the size of the bone graft. The passage of the graft in the growth plate or positioning of a fixation medium in the metaphysis can cause an arrest of the normal growth. The biological reasons are represented by possible complications of graft integration, and above all, by the thermal shock of mechanical drilling on the bone, which could cause heat necrosis and subsequently lead to epiphyseal fusion.<sup>15,16</sup>

The aim of this study was to verify the return of the skeletally immature professional athletes to sports in the long-term, following ACL reconstruction with the original all-inside technique and with manual drilling.

### Patients and methods

The design of our study is a single surgeon (GC), single technique, and retrospective series of ACL reconstruction employing the 'original all-inside technique'<sup>17,18</sup> on preadolescent patients with knee anterolateral rotatory instability after sustaining total tear of the ACL.

All reconstructions were performed by the senior author between the years 2002 and 2006. The Institutional Review Board approved the study.

Twenty-nine patients who had radiographic evidence of open physes, were less than 14 years of age at the time of surgery and those with a minimum follow-up period of eight years were included in the study. Twenty-four of them (14 boys, 10 girls; mean age: 13.15 years, range: 9–14 years) agreed to participate in the clinical interviews, examinations, and biomechanical and radiographic evaluations.

According to the Tanner scale, one patient was Tanner 1, three were Tanner 2, five were Tanner 3, twelve were Tanner 4 and one was Tanner 5 at the time of surgery. All patients sustained acute injuries and had no significant difference in limb length (<0.5 cm). Long-cassette radiographs were obtained preoperatively to measure the limb length, mechanical axis and the physeal length.

Fifteen of the patients were athletes in youth teams; 14 of them were footballers, four were volleyball players, three were basketball players, one was a cyclist, one was an artistic dancer, and one was a fighter.

Twelve patients had accompanying meniscal tears; six of them had longitudinal lesions of the medial meniscus, three had longitudinal lesions of the lateral meniscus and one had a bucket handle lesion of the medial meniscus. All these patients were surgically treated at the same time of the ACL reconstruction; two with selective meniscectomy and 10 with meniscal sutures using the Mulberry technique (three sutures failed and meniscectomy was performed as a revision surgery).

The original all-inside ACL reconstruction was performed under peripheral or general anesthesia and with the use of a tourniquet. Employing a lateral infrapatellar access and a standard anteromedial access, the chondral and meniscal lesions were treated when indicated. The notch was debrided and the remnants of the ACL were left when possible. 2.0-mm guide wires were introduced transphyseally into the femur and tibia with the out-in/free-hand

techniques and two small tunnels with a diameter of 4 mm were drilled both into the tibia and the femur (Fig. 1a and b).

Quadrupled gracilis tendon grafts were used for all ACL reconstructions. The grafts were harvested and measured in size, and the tibial and femoral half tunnels were manually drilled in-out in respect with the graft measurements (Fig. 2a and b). Then, the grafts were tensioned and fixed; femoral fixations were performed using the ENDOBUTTON polyester sutures and tibial fixations were carried out using small, resorbable interference screws from the outside.

Patients were immobilized in extension brace for 10 days after surgery. No weight-bearing was allowed for 21 days. Then, partial bearing was encouraged with two crutches for five days and with one crutch for four days. Full weight-bearing was allowed after 30 days. Isometric exercises were started as soon as possible and knee flexion was started on the 10th postoperative day. Playing sports was allowed only after biomechanical examinations confirmed a good muscular and articular functional recovery in isokinetic and kinematic tests. If the menisci were sutured, weight-bearing was forbidden for one month.

All recalled patients completed a questionnaire. The patients were facilitated by two blind examiners (MP and MB) who asked them to report their clinical and sportive histories. All patients completed a questionnaire, the IKDC subjective knee evaluation form and Tegner Activity Scale. One blinded biomechanics technician performed the biomechanical evaluations and one blinded examiner analyzed the biomechanical outcomes of the KT-1000




Fig. 1. (a, b) Introduction of guide wires into the femur and tibia with the out-in/free-hand techniques and drilling of the tunnels.

Download English Version:

<https://daneshyari.com/en/article/8795629>

Download Persian Version:

<https://daneshyari.com/article/8795629>

[Daneshyari.com](https://daneshyari.com)