

Contents lists available at [ScienceDirect](#)

Acta Orthopaedica et Traumatologica Turcica

journal homepage: <https://www.elsevier.com/locate/aott>

Regression of a symptomatic thoracic disc herniation with a calcified intervertebral disc component

Hidayet Sari ^a, Tugce Ozekli Misirlioglu ^{b,*}, Deniz Palamar ^a

^a Istanbul University, Cerrahpasa Medical Faculty, Physical Medicine and Rehabilitation Department, Istanbul, Turkey

^b Koc University Hospital, Physical Medicine and Rehabilitation Department, Istanbul, Turkey

ARTICLE INFO

Article history:

Received 19 September 2014

Received in revised form

22 February 2015

Accepted 17 May 2015

Available online xxx

Keywords:

Calcification

Pathologic

Herniated disc

Intervertebral disc

Thoracic vertebrae

ABSTRACT

There were only a few cases describing spontaneous regression of calcified thoracic disc herniation in the literature. We present a 38-year-old male office worker who had left paramedian-foraminal extruded disc at T7–T8 with calcifications of the T7–T8 and T8–T9 intervertebral discs. This case was unique in that the non-calcified extruded disc material regressed almost completely in 5 months while the calcified intervertebral discs remained the same during the process of regression. This report stresses that regression of the herniated material of the thoracic discs with subsidence of the symptoms is still possible even if the disc material is calcified.

© 2016 Turkish Association of Orthopaedics and Traumatology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Spontaneous regression or decrease in size of a herniated disc is commonly described in the lumbar and to a lesser extent in the cervical region while that of the thoracic region has rarely been reported.¹ In most of the cases, it was not indicated whether the discs that spontaneously regressed were calcified or not. There are only a few cases in the literature describing spontaneous regression of calcified thoracic disc herniation (TDH) in adults.^{2,3} We report a case of symptomatic TDH, in which the non-calcified extruded disc material regressed almost completely while the calcified intervertebral discs (IVDs) remained the same during the process of regression.

Case report

A 38-year-old male office worker, presented with a 2-month history of subscapular left-sided dorsal pain radiating in a belt-like distribution accompanied by occasional pain and paresthesia radiating to the left lower anterolateral chest wall. He reported that his pain had gradually increased and, recently had become unresponsive to medical treatment. His physical examination revealed

active myofascial trigger points (MTrP) in left lower paravertebral muscles in the thoracic spine with a predicted referred pain pattern compatible with the patient's symptoms. Upon diagnosing the patient with myofascial pain syndrome (MPS), we performed MTrP injections. He reported that his pain had decreased dramatically following the injection, but resumed a couple of days later. So, after we performed the MTrP injections two more times and observed the same partial relief, we re-evaluated the patient. Plain radiographs showed vague calcifications of the T7–T8 and T8–T9 IVDs with minimal degenerative changes of the corresponding end plates (Fig. 1A). Magnetic resonance imaging (MRI) revealed left paramedian-foraminal extruded disc at T7–T8 with left T7 nerve compression with hypointense signal in the central area of T7–T8 and T8–T9 discs (Fig. 2). We diagnosed the patient as having MPS accompanying T7–T8 TDH and, started a conservative treatment program including simple analgesic and non-steroid anti-inflammatory drugs with 15 sessions of physical therapy program. Gradually, his clinical symptoms subsided and, he returned to his activities of daily living without any complaints. Five months later, the patient was admitted to our clinics with his new X-ray and MRI scans that had been performed upon his request in a private hospital. We observed almost complete disappearance of the extruded fragment of the T7–T8 disc with a persisting hypointense zone in T8–T9 disc (Fig. 3). The hypointense zone in MRI was matched with the calcifications on X-ray (Fig. 1B) and, the patient was still asymptomatic.

* Corresponding author.

E-mail address: tozeklim@gmail.com (T.O. Misirlioglu).

Peer review under responsibility of Turkish Association of Orthopaedics and Traumatology.

<http://dx.doi.org/10.1016/j.aott.2015.05.002>

1017-995X/© 2016 Turkish Association of Orthopaedics and Traumatology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. (A) X-ray scan showing vague calcifications of the T7–T8 and T8–T9 IVDs; (B) 5 months later symptoms onset X-ray scan showing more prominent calcifications of the T7–T8 and T8–T9 IVDs.

Discussion

Thoracic disc herniations account for only about 0.25%–0.75% of all symptomatic herniated discs in adults.⁴ The diagnosis of symptomatic TDH is often overlooked because of the mismatch between the symptoms and location of the herniation.⁵ Disc calcification accompanying disc herniation is commonly reported in children and referred to as a ‘no touch lesion’ because of its benign nature and high rate of complete clinical and radiologic resolution.⁶ In adults, however, disc calcifications are thought to represent disc degeneration and herniation, and the rate of regression of calcified disc herniations are considered to be rare.³

Three mechanisms have been proposed to explain the process of disc regression. These are mechanical retraction of the herniated nucleus, gradual dehydration and shrinkage of the disc, and enzymatic degradation of herniated disc material.⁷ Among these, we find the third mechanism, namely, enzymatic degradation of herniated material, to be the most compelling and studied one. In that instance, extruded disc material is recognized as a foreign body and an inflammatory reaction is induced.

Almost 65% of herniated thoracic discs were shown to be calcified at presentation,⁸ and the presence of calcifications in the IVD is frequently found to be associated with hard herniated discs.⁹ Disc calcification is an important consideration since it is shown as



Fig. 2. (A) T2 weighted sagittal MRI scan showing the extruded disc at T7–T8 with hypointense signal in the central area of T7–T8 and T8–T9 discs; (B) T2 weighted axial MRI scan showing left paramedian-foraminal extruded disc at T7–T8 with left T7 nerve compression.

Download English Version:

<https://daneshyari.com/en/article/8795650>

Download Persian Version:

<https://daneshyari.com/article/8795650>

[Daneshyari.com](https://daneshyari.com)