

Accepted Manuscript

Title: Impact of motor fluctuations on real-life gait in Parkinson's patients

Authors: Ana Lgia Silva de Lima, Luc J.W. Evers, Tim Hahn, Nienke M. de Vries, Margaret Daeschler, Babak Boroogerdi, Dolors Terricabras, Max A. Little, Bastiaan R. Bloem, Marjan J. Faber



PII: S0966-6362(18)30283-2
DOI: <https://doi.org/10.1016/j.gaitpost.2018.03.045>
Reference: GAIPOS 6026

To appear in: *Gait & Posture*

Received date: 16-1-2018
Revised date: 26-3-2018
Accepted date: 27-3-2018

Please cite this article as: Silva de Lima Ana Lgia, Evers Luc JW, Hahn Tim, de Vries Nienke M, Daeschler Margaret, Boroogerdi Babak, Terricabras Dolors, Little Max A, Bloem Bastiaan R, Faber Marjan J. Impact of motor fluctuations on real-life gait in Parkinson's patients. *Gait and Posture* <https://doi.org/10.1016/j.gaitpost.2018.03.045>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Impact of motor fluctuations on real-life gait in Parkinson's patients

Impact of motor fluctuations on real-life gait in Parkinson's patients

Ana Lígia Silva de Lima^{a,b*}, Luc JW Evers^{a,c}, Tim Hahn^a, Nienke M de Vries^a, Margaret Daeschler^d,
Babak Boroojerdi^e, Dolores Terricabras^f, Max A. Little^{g,h}, Bastiaan R Bloem^a, Marjan J Faber^{a,i}

a – Radboud University Medical Center; Donders Institute for Brain, Cognition and Behaviour;
Department of Neurology; Nijmegen, The Netherlands

b – CAPES Foundation, Ministry of Education of Brazil, Brasília/DF, Brazil

c – Institute for Computing and Information Sciences, Nijmegen, The Netherlands

d – The Michael J Fox Foundation for Parkinson's Research, New York, USA

e – UCB Biopharma, Monheim, Germany

f – UCB Biopharma Slough, United Kingdom

g – Aston University, Birmingham, UK

h – Media Lab, Massachusetts Institute of Technology, Cambridge, USA

i – Radboud university medical center; Radboud Institute for Health Sciences; Scientific Center for
Quality of Healthcare; Nijmegen, the Netherlands

*Corresponding author: ana.silvadelima@radboudumc.nl. Radboud university medical center,
Department of Neurology, PO Box 9101 (920), 6500 HB Nijmegen, The Netherlands

Number of words: 2428; **Number of figures:** 3; **Number of tables:** 2

Highlights

- Parkinson's patients were more active between 8am and 1pm;
- Patients summed 72±39 (mean±standard deviation) minutes of walking per day.
- The severity of motor fluctuations did not correlate with time spent walking.
- Higher age and greater disease severity correlated with less time spent walking.
- The severity of motor fluctuations did not impact time spent walking after levodopa.

Download English Version:

<https://daneshyari.com/en/article/8798580>

Download Persian Version:

<https://daneshyari.com/article/8798580>

[Daneshyari.com](https://daneshyari.com)