


# Accepted Manuscript

Medial Overhang of the Tibial Component is associated with higher risk of inferior KOOS pain score after knee replacement.

C.S. Nielsen, A. Nebergall, J. Huddleston, T. Kallelose, H. Malchau, A. Troelsen



PII: S0883-5403(17)31133-6

DOI: [10.1016/j.arth.2017.12.027](https://doi.org/10.1016/j.arth.2017.12.027)

Reference: YARTH 56304

To appear in: *The Journal of Arthroplasty*

Received Date: 23 October 2017

Revised Date: 29 November 2017

Accepted Date: 15 December 2017

Please cite this article as: Nielsen C, Nebergall A, Huddleston J, Kallelose T, Malchau H, Troelsen A, Medial Overhang of the Tibial Component is associated with higher risk of inferior KOOS pain score after knee replacement., *The Journal of Arthroplasty* (2018), doi: 10.1016/j.arth.2017.12.027.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Title** Medial Overhang of the Tibial Component is associated with higher risk of inferior KOOS pain score after knee replacement.

**Authors:** Nielsen CS<sup>1,2</sup>, Nebergall A<sup>1</sup>, Huddleston J<sup>3</sup>, Kallemose T<sup>2</sup>, Malchau H<sup>1,4</sup>, Troelsen A<sup>2</sup>.

**Affiliations:**

<sup>1</sup>The Harris Orthopaedic Laboratory, Orthopaedic Department, Massachusetts General Hospital, Boston, USA;

<sup>2</sup>Department of Orthopaedic surgery, Copenhagen University Hospital Hvidovre, Copenhagen, Denmark,

<sup>3</sup>Department of Orthopaedic Surgery, Stanford University Medical Center, Stanford, California, USA, <sup>4</sup>L Sahlgrenska University Hospital, Department of Orthopaedics, Gothenburg, Sweden.

**Pages of manuscript: 9**

**Blinded information, page 3, line 56-58 in manuscript:**

112 patients were enrolled in Europe from 3 centers (2 centers in Denmark and 1 in Sweden); 98 patients in the United States from 4 centers (Massachusetts, California, Wisconsin and Tennessee), 81 patients from Asia all from 2 South Korean centers, and 32 patients from 1 center in Australia (Sydney).

Download English Version:

<https://daneshyari.com/en/article/8799369>

Download Persian Version:

<https://daneshyari.com/article/8799369>

[Daneshyari.com](https://daneshyari.com)