

Accepted Manuscript

Title: Influence of treatment modality on morbidity and mortality in periprosthetic femoral fracture. A comparative study of 71 fractures treated by internal fixation or femoral implant revision.

Author: S. Cohen X. Flecher S. Parratte M. Ollivier J.N. Argenson

PII: S1877-0568(18)30043-4
DOI: <https://doi.org/doi:10.1016/j.otsr.2017.12.018>
Reference: OTSR 1956

To appear in:

Received date: 7-6-2016
Accepted date: 1-12-2017

Please cite this article as: Cohen S, Flecher X, Parratte S, Ollivier M, Argenson JN, Influence of treatment modality on morbidity and mortality in periprosthetic femoral fracture. A comparative study of 71 fractures treated by internal fixation or femoral implant revision., *Orthopaedics and Traumatology: Surgery and Research* (2018), <https://doi.org/10.1016/j.otsr.2017.12.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Original article

Influence of treatment modality on morbidity and mortality in periprosthetic femoral fracture. A comparative study of 71 fractures treated by internal fixation or femoral implant revision.

S. Cohen^a, X. Flecher^{a,*}, S. Parratte^a, M. Ollivier^a, J.-N. Argenson^a

^a Institut du Mouvement et de l'Appareil Locomoteur, UMR CNRS 7287 /AMU, Centre Hospitalo-Universitaire Sud, Hôpital Sainte Marguerite, Boulevard de Sainte Marguerite, 13009 Marseille, France

* Corresponding author: Xavier Flecher, Service de Chirurgie Orthopédique, Hôpital Sainte Marguerite, Boulevard de Sainte Marguerite, 13009 Marseille, France

Tel.: (+33)(0)491745011

Fax: (+33)(0)491745625

E-mail: xavier.flecher@gmail.com

Abstract

Introduction

The incidence of periprosthetic femoral fracture is constantly increasing, with high associated morbidity and mortality. Surgical treatment is guided by the Vancouver classification, but the influence of type of treatment on morbidity and mortality has been

Download English Version:

<https://daneshyari.com/en/article/8802205>

Download Persian Version:

<https://daneshyari.com/article/8802205>

[Daneshyari.com](https://daneshyari.com)