

# Accepted Manuscript

Title: Application of neurite orientation dispersion and density imaging or diffusion tensor imaging to quantify the severity of cervical spondylotic myelopathy and assess postoperative neurological recovery

Author: Genki Okita, Tetsuro Ohba, Tomohiro Takamura, Shigeto Ebata, Ryo Ueda, Hiroshi Onishi, Hirotaka Haro, Masaaki Hori



PII: S1529-9430(17)30325-X  
DOI: <http://dx.doi.org/doi: 10.1016/j.spinee.2017.07.007>  
Reference: SPINEE 57391

To appear in: *The Spine Journal*

Received date: 27-2-2017  
Revised date: 30-5-2017  
Accepted date: 6-7-2017

Please cite this article as: Genki Okita, Tetsuro Ohba, Tomohiro Takamura, Shigeto Ebata, Ryo Ueda, Hiroshi Onishi, Hirotaka Haro, Masaaki Hori, Application of neurite orientation dispersion and density imaging or diffusion tensor imaging to quantify the severity of cervical spondylotic myelopathy and assess postoperative neurological recovery, *The Spine Journal* (2017), <http://dx.doi.org/doi: 10.1016/j.spinee.2017.07.007>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Application of neurite orientation dispersion and density imaging or diffusion**  
2 **tensor imaging to quantify the severity of cervical spondylotic myelopathy and**  
3 **assess postoperative neurological recovery**

4

5 Genki Okita, MD,<sup>1</sup> Tetsuro Ohba, MD, PhD,<sup>1\*</sup> Tomohiro Takamura, MD, PhD,<sup>2</sup> Shigeto

6 Ebata, MD, PhD,<sup>1</sup> Ryo Ueda, BS,<sup>3,4</sup> Hiroshi Onishi, MD, PhD,<sup>2</sup> Hirotaka Haro, MD,

7 PhD,<sup>1</sup> Masaaki Hori, MD, PhD<sup>2,3</sup>

8 <sup>1</sup>Department of Orthopaedic Surgery, University of Yamanashi, Japan

9 <sup>2</sup>Department of Radiology, University of Yamanashi, Japan

10 <sup>3</sup>Department of Radiology, School of Medicine, Juntendo University, Japan

11 <sup>4</sup>Department of Radiological Science, Tokyo Metropolitan University of Health  
12 Sciences, Japan

13 \*Address correspondence and reprint requests to:

14 Tetsuro Ohba, MD, PhD

15 Department of Orthopaedics, University of Yamanashi

16 1110, Shimokato, Chuo, Yamanashi 409-3898, Japan

17 E-mail: tooba@yamanashi.ac.jp

18 Phone: +81-55-273-6768; Fax: +81-55-273-9241

19 No financial or material support was obtained for this study. There is no conflict of  
20 interest.

21 **Acknowledgments**

22 This work was supported by JSPS KAKENHI Grant Number 16K10328. The authors  
23 thank Hiroshi Kumagai for his cooperation with the imaging technique

24

25

26

Download English Version:

<https://daneshyari.com/en/article/8804569>

Download Persian Version:

<https://daneshyari.com/article/8804569>

[Daneshyari.com](https://daneshyari.com)