



Egyptian Society of Ear, Nose, Throat and Allied Sciences  
**Egyptian Journal of Ear, Nose, Throat and Allied Sciences**

[www.ejentas.com](http://www.ejentas.com)



ORIGINAL ARTICLE

# Role of endoscopic guided partial adenoidectomy in avoiding open nasality



Eman Mostafa \*, Ibrahim Rezk

Sohag University Hospital, Egypt

Received 3 March 2016; accepted 20 June 2016

## KEYWORDS

Partial endoscopic adenoidectomy;  
Resonance of speech;  
Nasometry

**Abstract** *Objectives:* Assess the effects of endoscopic guided partial adenoidectomy on resonance of speech in patients with poor palatal mobility and evaluate its role in avoiding postoperative open nasality.

*Materials and methods:* This study was a prospective randomized trial conducted in one year duration. A total of 40 patients were assessed for eligibility; 24 patients were excluded as they did not meet the inclusion criteria. Two patients out of the eligible sixteen refused surgery. Inclusion criteria: 1-patient complaining of snoring, nasal obstruction and nasal discharge, all of which were caused by adenoid. 2-Poor palatal mobility on endoscopic examination. Exclusion criteria: any neurological deficit, muscular disorder or structural defects of the palate. All eligible patients had undergone partial adenoidectomy. Speech was evaluated preoperative and postoperative.

*Results:* The study group had definite poor palatal mobility on endoscopic examination and lateral videofluoroscopy. Endoscopic guided partial adenoidectomy has been done to all 14 patients. Postoperative evaluation showed marked relieving of symptoms such as nasal obstruction, discharge and mouth breathing with successfully maintaining the velopharyngeal competence but there was no complete relieving of closed nasality.

*Conclusion:* There is necessity to screen potential candidates for adenoidectomy in order to prevent postoperative velopharyngeal dysfunction. Nasoendoscopy and lateral videofluoroscopy have been useful to examine the palatal mobility, palatal length, depth of the pharynx and to exclude possible occult cleft palate. Partial adenoidectomy should be taken into consideration for risky patients. Satisfactory results were obtained as regards relieving nasal symptoms and maintaining velopharyngeal competence but didn't improve quite fully the closed nasality.

© 2016 Egyptian Society of Ear, Nose, Throat and Allied Sciences. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Sound energy begins when the vocal folds vibrate, producing sound. The sound energy travels in a superior direction through a series of interconnected resonators that include the pharynx, the oral cavity, and the nasal cavity. The size and

\* Corresponding author.

E-mail address: [eman\\_mostafa@med.sohag.edu.eg](mailto:eman_mostafa@med.sohag.edu.eg) (E. Mostafa).

Peer review under responsibility of Egyptian Society of Ear, Nose, Throat and Allied Sciences.

<http://dx.doi.org/10.1016/j.ejenta.2016.06.003>

2090-0740 © 2016 Egyptian Society of Ear, Nose, Throat and Allied Sciences. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

shape of the resonating cavities directly affect the perceived resonance and voice quality. The velopharyngeal mechanism is responsible for regulating and directing the transmission of sound energy and air pressure in the oral and nasal cavities.<sup>1</sup> For normal speech and resonance, velopharyngeal closure should be complete during the production of oral sounds; and for nasal sounds, sound energy should be relatively unimpeded through the pharynx and nasal cavity.<sup>2</sup>

Closed nasality is a reduction in nasal resonance during phonation, especially of nasal phonemes such as /m/, /n/, and /ng/. It typically results from either partial or complete blockage of the nasal cavity or nasopharynx from mucosal edema associated with viral upper respiratory infection (URI), hypertrophic tonsils/adenoids, allergic rhinitis, sinusitis, hypertrophic turbinates, or anatomic obstruction from a deviated nasal septum or choanal atresia.<sup>3</sup>

In 1958, Gibb<sup>4</sup> indicated an incidence of hypernasality postadenoidectomy in approximately 1 of 2000 cases. Closure pattern of velopharyngeal valve in typical children is veloade-noidal rather than velopharyngeal closure.<sup>5</sup> Adenoid mass is vital to velopharyngeal closure in children and removal necessitates a change in the pattern of velopharyngeal valving.<sup>6</sup> These changes are easily overcome if there is no anatomic abnormality.

The aim of our study was to assess the effects of endoscopic guided partial adenoidectomy on resonance of speech and evaluate its role in avoiding postoperative open nasality.

## 2. Material

This study was a prospective randomized trial conducted for a one year duration in Sohag University Hospital (SUH). A total of 40 patients were assessed for eligibility age of the patients ranged from 4 to 21 years. Inclusion criteria: 1-patient complaining of snoring, nasal obstruction and nasal discharge, all of which were caused by adenoid. 2-Poor palatal mobility on endoscopic examination. Exclusion criteria: any neurological deficit, muscular disorder or structural defects of the palate such as cleft palate or submucous cleft palate. Twenty four patients were excluded as they had good palatal mobility on endoscopic examination. Two patients out of the eligible sixteen refused surgery.


Approval of the ethics committee of Sohag University was obtained. A written consent to participate and to publish was taken from all patients or participants before our study procedures. None of the authors have any competing interests.

All patients had undergone ear, nose and throat examination. They have been referred to the Phoniatrics Unit in Sohag University hospital for evaluation of the auditory perceptual assessment to assess voice and speech as regards resonance (open or closed or mixed nasality). Perceptual speech assessment involves the evaluation of all the components of speech production (including voice), articulation (including misarticulations, motor speech, oral motor sequencing), velopharyngeal function, and overall speech intelligibility.

Palatal mobility was assessed through oral examination. The relative length of the velum as well as its mobility during phonation was assessed. The velum should be raised and the velar “dimple” should be back approximately 80% of the length of the soft palate. Nasoendoscopic examination was done for further evaluation of the palatal mobility. Nasoendoscopy provides a useful tool for direct visualization of the VP during speech, evaluating the velopharyngeal function and allowing for a 2-dimensional view of the VP complex. This allows the physician to evaluate lateral wall and palatal mobility during speech; determine the presence of a notch on the nasal surface of the palate; the Passavant ridge and aberrant pulsations; and estimate gap size and shape if present. In normal velopharyngeal closure, there should be good sphincter function and optimum closure during phonation.

Videofluoroscopy was also done in selected patients (4 cases) when needed. A videofluoroscopy speech study is a radiographic evaluation that allows the direct visualization of all aspects of the velopharyngeal sphincter during speech. It was done in selected cases such as uncooperative child during nasoendoscopy, excessive mucous, the need to evaluate the thickness and length of the palate. Videofluoroscopy is used as a complementary tool to nasoendoscopy in evaluating palatal mobility as a protocol that is followed in SUH. In both procedures, the patient is asked to repeat standard vowels, consonants and sentences so that the palatal mobility is evaluated (Fig. 1).

Endoscopic guided partial adenoidectomy using the powered microdebrider was done for the risky patients with mild poor palatal mobility on endoscopic examination ( $n = 14$ ). All eligible patients had undergone partial adenoidectomy only with no tonsillectomy. Partial adenoidectomy was started high in the nasopharynx from upper limit of adenoid tissue, choanal parts. Resection was continued in side to side fashion sparing the lower half of the adenoid.<sup>7</sup> The cutting and aspirating action of the shaver and simultaneous irrigation removes both adenoid tissue and the blood, thus providing a clear view. Better control of the depth of removal of adenoid is achieved thus



**Figure 1** (A) Lateral videofluoroscopy at rest. (B) Lateral videofluoroscopy while saying the vowel /a/.

Download English Version:

<https://daneshyari.com/en/article/8805803>

Download Persian Version:

<https://daneshyari.com/article/8805803>

[Daneshyari.com](https://daneshyari.com)