

HOSTED BY



Contents lists available at ScienceDirect

International Journal of Pediatrics and Adolescent Medicine

journal homepage: <http://www.elsevier.com/locate/ijpam>

Disc battery ingestion; a single event with different outcomes

E. Sindi ^a, A. Bahadur ^b, H. AlRajhi ^b, Y. Yousef ^{a, b, c, *}^a King AbdulAziz Medical City (KAMC), Jeddah, Saudi Arabia^b King Saud Bin AbdulAziz University for Health Sciences, College of Medicine- Jeddah (KSAU-HS, COM-J), Jeddah, Saudi Arabia^c King Abdullah International Medical Research Center (KAIMRC), Jeddah, Saudi Arabia

ARTICLE INFO

Article history:

Received 9 March 2017

Received in revised form

24 April 2017

Accepted 25 April 2017

Available online xxx

Keywords:

Disc battery

Foreign body ingestion

Acquired tracheoesophageal fistula

Esophageal stricture

ABSTRACT

Foreign body (FB) ingestion is a common problem especially in children below the age of 5 years. This is fueled by their curiosity to explore their surroundings. The ingested foreign body finds its way out of the gastrointestinal tract without any serious consequences most of the time. On the other hand, disc battery ingestion has been reported to cause serious harm when ingested including death. We report two patients who had ingested disc batteries and their respective outcomes.

© 2017 Publishing services provided by Elsevier B.V. on behalf of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Infants and young children are known to explore their surroundings by touch and taste. It is part of achieving their developmental milestones. Many objects find their way to a child's gastrointestinal tract and usually these objects find their way out. Disc batteries have become more common in our homes and so they have become part of a child's surroundings. As disc batteries are smooth and shiny, they are attractive to this age category. Therefore, ingestion of disc battery is becoming a more frequent encounter. Complications of disc battery ingestion are many, ranging from dysphagia to more severe outcomes like esophageal burns and tracheoesophageal fistula. Most complications can be minimized by early detection and intervention. Application of preventive measures are needed if we want to eliminate this event. We are reporting two different patients who had disc battery ingestion and their respective outcomes. Also, we included a background literature review.

1.1. Case 1

A previously healthy 2-year-old girl presented to our emergency department (ED) after her mother noticed that she was drooling for 2 days. The mother told us the child was interested in disc batteries and she had repeatedly found her playing with them. The child was previously seen at a polyclinic where she was diagnosed with pharyngitis, for which she was given antibiotics. Next day, she developed fever and vomiting. A plain chest x-ray confirmed that the patient had a disc battery in the upper esophagus. The parents were advised to take their daughter to a hospital.

Upon presentation to our (ED), she had shortness of breath, cough, and drooling for the past 48 hours. She was ill-looking with a heart rate of 150 beats per minute, blood pressure of 99/56 mmHg, respiratory rate of 34 breaths per minute, and a temperature of 40 degrees Celsius (°C). A Chest x-ray showed a round disc-like opacity projecting over the esophagus. Both costophrenic angles were clear. The cardiac shadow was within normal limits. No pneumonia or pneumothorax was noted (Fig. 1a and b).

The patient was admitted and taken to the operating room for urgent esophagoscopy and retrieval of the foreign body. Esophagoscopy revealed circumferential burn at the site of the disc battery with moderate to severe inflammatory changes in the mucosa. The disc battery was removed very carefully. The esophagus was inspected again; no signs of perforation were seen and the procedure was ended without advancing the scope beyond the area of the burn. The patient was shifted, ventilated and intubated to the

* Corresponding author. King AbdulAziz Medical City (KAMC), Jeddah, Saudi Arabia.

E-mail addresses: enassindi@gmail.com (E. Sindi), ay.ab777@gmail.com (A. Bahadur), dr.hamdan.ra@hotmail.com (H. AlRajhi), yousefyfya@ngha.med.sa, yamyousef@gmail.com (Y. Yousef).

Peer review under responsibility of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia.

<http://dx.doi.org/10.1016/j.ijpam.2017.04.003>

2352-6467/© 2017 Publishing services provided by Elsevier B.V. on behalf of King Faisal Specialist Hospital & Research Centre (General Organization), Saudi Arabia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. **a** PA chest xray showing a round disc battery in the upper esophagus. No pneumomediastinum or signs of pneumonia noted. **b** Lateral chest xray showing disc battery to be in the esophagus not the trachea. Trachia is anterior to The Foreign Body.

pediatric intensive care unit (PICU) for observation for 24 hours. A post-operative chest x-ray revealed no pneumomediastinum. So, she recovered slowly and extubated. On the ward, the patient was kept nil per mouth. She received intravenous (IV) hydration, histamine receptors blocker (H2 blocker) and amoxicillin/clavulanic acid IV for five days. Later; she had a water-soluble contrast swallow which showed no leak and no stricture in the esophagus. After which, she was started on clear fluids and progressed gradually to soft diet. The patient was discharged home on ranitidine (2mg/kg) per 24 hours in two divided doses. Instructions were given to come to the ED in case of fever, drooling or difficulty swallowing.

The patient was seen in the clinic after one month. She was tolerating her food well. A repeat esophagogram did not show any stricture. Ranitidine was stopped. The patient remained well on repeated outpatient follow up 3 and 6 months later.

1.2. Case 2

A 15 months old girl presented to another hospital's ED due to a history of difficulty swallowing that started one week before presentation. Chest x-ray revealed a round disc like foreign body in the esophagus. The patient was admitted there and an upper gastrointestinal (GI) flexible endoscopy was done. It revealed a disc battery impacted in the upper part of the esophagus which was removed. During the procedure, a very deep large esophageal ulceration with gangrenous tissue and local bleeding was found possibly due to long-lasting impaction.

The patient was admitted to PICU for five days due to stridor which started on the day of presentation to the hospital. On the sixth day, she was transferred to the pediatric floor on nasal cannula oxygen therapy, IV antibiotics, total parenteral nutrition (TPN), and frequent suctioning from the mouth due to excess secretions.

While on the ward, the patient suffered three attacks of choking and cyanosis which improved on suctioning and oxygen. Examination at that time, showed the patient to be fully conscious, having stable vital signs, with a lot of mouth secretions. Chest examination showed equal bilateral air entry with transmitted sounds and coarse crepitation. Complete blood count (CBC), C-reactive protein (CRP), renal panel, bone biochemistry, and blood culture were all within normal range.

Computed tomography (CT) scan of the chest with oral contrast showed no evidence of pneumomediastinum. The contrast was

seen around the esophagus middle and lower parts mostly in keeping with periesophageal leak. No significant amount of contrast was noted inside the trachea and its main branches, tracheoesophageal fistula possibility was raised.

The patient was kept under close observation, respiratory care, TPN and antibiotics (Meropenem, vancomycin) for 6 weeks.

Afterwards, she underwent diagnostic esophagoscopy which proved the presence of the fistula. Repair of tracheoesophageal fistula with pericardial patch interposition through a right thoracotomy was done.

The first contrast study after the repair showed no stricture or leak. The child was discharged home on anti-reflux medication.

The patient initially presented to us about 5 weeks after tracheoesophageal fistula (TEF) repair. She was only allowed to take fluid diet since the day of surgery. On examination, she was healthy and afebrile with clean right thoracotomy wound. An upper GI contrast study showed mild narrowing but good flow of contrast at the repair site with no evidence of stricture. She was allowed full diet with regular follow-up appointments with our outpatient service.

Three months after the repair, the mother reported noticing difficulty swallowing solid foods, choking and some drooling as well. An urgent contrast study (Fig. 2a and b) showed a very tight esophageal stricture.

The patient was admitted on semi urgent basis for esophageal dilatation. Esophagoscopy and attempted esophageal dilatation failed due to failure of passing a guidewire into the stomach. There were multiple foreign bodies that were seen which looked like peanut pieces. Those were retrieved using the optical forceps. Upon retrieval of these foreign bodies, there was some inflammation and bleeding at the esophagus where the foreign bodies were impacted so the procedure was stopped. The patient was given steroids and antibiotics for several days. She was taken again to the theater 4 days later. This time, ureteric dilators were used followed by balloon dilation up to the size 10mm. The patient tolerated the procedure well and recovered uneventfully. She tolerated a soft diet and was discharged on Ranitidine (2mg/kg/day) with an elective readmission for another dilatation in one month.

The patient's second dilatation went smoothly. Savory Gillard dilators were used up to size of 12mm. Six weeks later, the third dilatation was done with reaching up to the size of 12.8mm easily. The last dilatation was done after another 6-weeks interval.

Download English Version:

<https://daneshyari.com/en/article/8809626>

Download Persian Version:

<https://daneshyari.com/article/8809626>

[Daneshyari.com](https://daneshyari.com)