



Contents lists available at ScienceDirect

Journal of Pediatric Surgery

journal homepage: www.elsevier.com/locate/jped surg

Morbidity and mortality among “big” babies who develop necrotizing enterocolitis: A prospective multicenter cohort analysis

Cristine S. Velazco^a, Brenna S. Fullerton^a, Charles R. Hong^a, Kate A. Morrow^c, Erika M. Edwards^{b,c}, Roger F. Soll^{b,c}, Tom Jaksic^a, Jeffrey D. Horbar^{b,c}, Biren P. Modi^{a,*}

^a Center for Advanced Intestinal Rehabilitation, Department of Surgery, Boston Children's Hospital and Harvard Medical School, Boston, MA

^b University of Vermont, Burlington, VT

^c Vermont Oxford Network, Burlington, VT

ARTICLE INFO

Article history:

Received 12 September 2017

Accepted 5 October 2017

Available online xxx

Key words:

Necrotizing enterocolitis

Congenital anomalies

Growth

Surgery

ABSTRACT

Background: Necrotizing enterocolitis (NEC) is classically a disease of prematurity, with less reported regarding morbidity and mortality of this disease among other infants.

Methods: Data were prospectively collected from 2009 to 2015 at 252 Vermont Oxford Network member centers on neonates with birth weight > 2500 g admitted to a participating NICU within 28 days of birth.

Results: Of 1629 neonates with NEC, gestational age was 37 (36, 39) weeks, and 45% had major congenital anomalies, most commonly gastrointestinal defects (20%), congenital heart defects (18%), and chromosomal anomalies (7%). For the 23% of infants who had surgery for NEC, mortality and length of stay were 23% and 63 (36, 94) days versus 8% and 34 (22, 61) days in medical NEC. Independent predictors of mortality were congenital heart defects ($p < 0.0001$), chromosomal abnormalities ($p < 0.05$), other congenital malformations ($p < 0.001$), surgical NEC ($p < 0.0001$), and sepsis ($p < 0.05$). All of these in addition to gastrointestinal defects were independent predictors of increased length of stay. Nutritional morbidity at discharge included 6% receiving no enteral feeds and 27% who were <10th percentile weight-for-age.

Conclusions: Major congenital anomalies are present in nearly half of >2500 g birth weight infants diagnosed with necrotizing enterocolitis. Morbidity and mortality increase with sepsis, surgical disease, and congenital anomalies.

Type of study: Prognosis Study.

Level of evidence: Level II.

© 2017 Elsevier Inc. All rights reserved.

Necrotizing enterocolitis (NEC) is the most common surgical emergency in neonates and is classically a disease of prematurity [1], with approximately 90% of cases diagnosed in preterm infants [2]. Among these preterm infants, the incidence and mortality of NEC decrease with higher birth weights [3]. While NEC has been extensively described in preterm infants, less is known about the heterogeneous group of normal birth weight infants who are diagnosed with NEC. In full-term infants, NEC is often associated with comorbidities where factors such as asphyxia or metabolic abnormalities may be at play; for example, congenital heart disease is associated with increased risk for NEC in both VLBW and full-term infants [2,4]. We utilized a prospectively collected quality improvement database from a collaborative of neonatal intensive care units to: 1) obtain benchmark data on mortality and length of stay among neonates with birth weight > 2500 g who were diagnosed with NEC, 2) identify independent risk factors for these outcomes and 3) explore other potential morbidities in this patient population.

1. Method

1.1. Study designs

Vermont Oxford Network (VON) is a nonprofit voluntary collaboration of health professionals dedicated to improving the care of high risk infants and their families (www.vtoxford.org). In support of this mission, VON maintains databases for use in quality improvement. Members may report data for VLBW infants only (defined as either a gestational age of 22 to 29 weeks or a birth weight of 401 to 1500 g) or report data for all NICU admissions regardless of birth weight or gestational age. Between 2009 and 2015, 1074 hospitals participated in VON with 489 reporting data on all NICU infants. Data are centrally collected and analyzed, and strict definitions are applied using a universal Manual of Operations.

Infants at participating North American centers diagnosed with NEC who had birth weights greater than 2500 g were included in this study. One Canadian center and 251 American centers contributed data for at least one included infant (Appendix 1; online only). Survival was defined as hospitalization at 1 year of age or discharge home without readmission.

* Corresponding author at: Boston Children's Hospital and Harvard Medical School, 300 Longwood Avenue, Fegan 3, Boston, MA 02115. Tel.: +1 617 355 9600; fax: +1 617 730 0477.

E-mail address: biren.modi@childrens.harvard.edu (B.P. Modi).

Clinical diagnosis of NEC was defined by at least one physical finding (bilious gastric aspirate or emesis, abdominal distention, or occult/gross blood in the stool in the absence of anal fissures) and at least one radiographic finding (pneumatosis intestinalis, hepatobiliary gas, or pneumoperitoneum). NEC could also be diagnosed at laparotomy or at postmortem examination. For each infant diagnosed with NEC, type of surgery for NEC was recorded, if applicable. For each infant, all surgical procedures were reported from a list of defined procedures in the VON Manual of Operations [5].

Any major congenital anomalies, as defined by the VON Manual of Operations, were reported for each infant [5], with some infants having multiple anomalies. All reported anomalies not specifically defined by an available code were reviewed by surgeons and appropriately classified. Isolated atrial or ventricular septal defects were not included as serious congenital heart disease, as no data on clinical severity were collected.

Sepsis was defined by cerebral spinal fluid or blood culture positive for fungus or any of 30 defined bacterial pathogens. Cultures positive for coagulase negative staphylococcus were only included if the infant also had signs of generalized infection and underwent treatment with five or more days of intravenous antibiotics after the culture was obtained. Weight for age percentile was calculated using the Fenton growth standards for all infants discharged home at less than 50 weeks post menstrual age [6], and using the World Health Organization growth standards adjusted for gestational age for all infants discharged home at greater than 50 weeks post menstrual age [7]. Infants with a less than 15 day length of stay were not included in calculations of weight for age at discharge, so that physiological neonatal weight loss did not bias the results [8].

“Initial resuscitation” refers to interventions performed in the delivery room or in a resuscitation area immediately following birth and prior to NICU admission. Resuscitation strategies recorded include the use of supplemental oxygen, face mask ventilation, nasal continuous positive airway pressure (CPAP), endotracheal tube ventilation, the use of epinephrine, and cardiac compressions. Assisted ventilation was defined as intermittent positive pressure ventilation through an endotracheal tube with a conventional ventilator (intermittent mandatory ventilation rate < 240/min) at any time after leaving the initial resuscitation area. The number of days of assisted ventilation includes any complete or partial day during which the infant received ventilation via endotracheal intubation and a conventional ventilator.

This study was performed as part of an ongoing collaboration between VON and pediatric surgeons at Boston Children’s Hospital. The Committee on Human Research at the University of Vermont approved the use of the Vermont Oxford Network Research Repository for this analysis (#15-143), and IRB exemption was granted at Boston Children’s Hospital (IRB-P00002185).

1.2. Statistical methods

Multivariate logistic regression was used to assess risk factors for mortality and Poisson regression was used to assess risk factors for increased length of stay (LOS) among surviving infants, controlling for clustering of infants within hospitals. Models included all variables in Table 1; those with P-values < 0.05 were considered independent predictors. All analyses were produced using SAS version 9.4 (SAS Institute, Cary, NC).

2. Results

There were 252 centers that contributed data on 1629 neonates > 2500 g birth weight (60% male) who developed NEC. Median (IQR) gestational age in this cohort was 37 (36, 39) weeks (Table 1). Major congenital anomalies were seen in 44%, most commonly gastrointestinal defects (20%), congenital heart defects (18%), and chromosomal anomalies (7%); 10% had other congenital anomalies.

Of the 1629 included infants with NEC, 380 (23%) underwent surgery for NEC (surgical NEC), including 39 infants who received primary

Table 1

Patient characteristics, comparing infants with medical and surgical NEC.

	Medical NEC	Surgical NEC
Number of infants	1249	380
Birth Weight (g: median, Q1, Q3)	3035 (2754, 3453)	3010 (2739, 3405)
Gestational Age (weeks; median, Q1, Q3)	37 (36, 39)	37 (36, 39)
Apgar		
1 min (median, Q1, Q3)	8 (6, 8)	8 (5, 8)
5 min (median, Q1, Q3)	9 (8, 9)	9 (8, 9)
Male (%)	60	60
Maternal Hypertension (%)	15	21
Chorioamnionitis (%)	5	6
Small for Gestational Age (%)	8	10
Multiple Gestation (%)	4	2
Cesarean Delivery (%)	45	46
Outborn (%)	46	58
Major Birth Defect (%)	42	51
Any Initial Resuscitation (%)	52	50
Oxygen	49	48
Face mask	25	26
Intubation	13	18
CPAP	14	14
Sepsis (%)	14	27
Respiratory Distress Syndrome (%)	14	12
Assisted Ventilation (%)	59	97
Duration of Assisted Ventilation (median, Q1, Q3)	1 (0, 7)	7 (3, 18)
Any abdominal surgery (%)	35	100

NEC: necrotizing enterocolitis. CPAP: continuous positive airway pressure.

peritoneal drainage (PPD). Of the infants who received PPD, 30 also underwent laparotomy (with 7 deaths in this group), five died without further surgical intervention, and four survived to discharge without any additional abdominal surgeries. Mortality was 22% among infants with surgical NEC and 8% among infants with medical NEC (ARR 2.2 (1.7–2.9), with median (IQR) length of stay among survivors of 63 (36, 94) and 34 (22, 61) days, respectively ($p < 0.0001$). The proportion of infants with surgical NEC who had a major birth defect was 51% compared to 42% of those infants with medical NEC. Fig. 1 demonstrates prevalence and percent mortality of each significant independent predictor of death identified; the top of each gray bar indicates the proportion of infants in this study affected by each given risk factor, which ranged from 7% for chromosomal anomalies to 23% for surgical NEC. Infants who survived are represented by the light gray portion of the bar, and nonsurvivors are represented by the dark gray portion, with percent mortality for infants with each risk factor displayed.

Overall mortality for NEC was 11% in this cohort, and 5% among infants without any major congenital anomaly. Significant independent predictors of mortality on multivariate logistic regression were congenital heart defects ($p < 0.0001$), chromosomal abnormalities ($p < 0.05$), other congenital malformations ($p < 0.001$), surgical NEC ($p < 0.0001$), and sepsis ($p < 0.05$) (Table 2). Risk factors for increased length of stay among survivors included the risk factors for mortality, listed above, and the presence of gastrointestinal defects ($p < 0.0001$) (Table 3).

In exploring additional outcomes, major nutritional morbidity was observed among survivors. Despite a low prevalence of small for gestational age infants, as would be expected in this cohort of infants selected by birth weight criteria, by the time of discharge the percent of infants who were less than the tenth percentile weight for age had increased (Table 4). At the time of discharge, 7% of the infants with medical NEC were receiving no enteral feeds, and 3% of those with surgical NEC were receiving no enteral feeds. During the initial hospital stay, 14% of infants underwent surgery for gastrostomy or jejunostomy tube placement. Data on the percent of infants discharged on a combination of enteral and parenteral support are not available. At discharge, approximately one quarter of the cohort were receiving exclusively formula feeds (24% for medical NEC, 27% for surgical NEC), half were

Download English Version:

<https://daneshyari.com/en/article/8810576>

Download Persian Version:

<https://daneshyari.com/article/8810576>

[Daneshyari.com](https://daneshyari.com)