


Accepted Manuscript

The role of pre operative intra muscular testosterone in improving functional and cosmetic outcomes following hypospadias repair: A prospective randomized study

Ramesh Babu, Sibi Chakravarthi



PII: S1477-5131(17)30300-5

DOI: [10.1016/j.jpurol.2017.07.009](https://doi.org/10.1016/j.jpurol.2017.07.009)

Reference: JPUROL 2616

To appear in: *Journal of Pediatric Urology*

Received Date: 20 May 2017

Revised Date: 1477-5131 1477-5131

Accepted Date: 3 July 2017

Please cite this article as: Babu R, Chakravarthi S, The role of pre operative intra muscular testosterone in improving functional and cosmetic outcomes following hypospadias repair: A prospective randomized study, *Journal of Pediatric Urology* (2017), doi: 10.1016/j.jpurol.2017.07.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The role of preoperative intramuscular testosterone in improving functional and cosmetic outcomes following hypospadias repair: A prospective randomized study

Ramesh Babu *, Sibi Chakravarthi

Department of Pediatric Urology, Sri Ramachandra Medical College and Research Institute, Porur, Chennai, India

* **Corresponding author.** Pediatric Urology Unit, Sri Ramachandra Medical College and Research Institute, Porur, Chennai 600116, India.

E-mail address: drrameshbabu1@gmail.com (R. Babu)

Summary

Background

There are very few well-designed studies on the role of preoperative testosterone in hypospadias repair. The aims of the current study are to ascertain the efficacy of intramuscular testosterone in increasing the glans width, reducing the complications and improving the functional and cosmetic outcomes.

Materials and methods

Between January 2013 and December 2016, a total of 200 patients with distal hypospadias were recruited to this prospective randomized study. Only patients with distal hypospadias eligible for a tubularized incised plate (TIP) repair were included. Group 1 did not receive any intervention, and group 2 received three doses of intramuscular testosterone enanthate 2 mg/kg at the age of 9, 10, and 11 months. Glans width was measured in millimeters in all patients using calipers at the time of recruitment and at the time of surgery. Within group 2, those who failed to have more than a 2 mm increase in size were considered non-responders (group 2a), and the remaining were taken to be responders (group 2b). TIP repair was performed by the same surgeon, and patients were followed

Download English Version:

<https://daneshyari.com/en/article/8811712>

Download Persian Version:

<https://daneshyari.com/article/8811712>

[Daneshyari.com](https://daneshyari.com)