


Accepted Manuscript

The incidence of isolated penile torsion in north India: A study of 5,018 male neonates

Amilal Bhat, Mahakshit Bhat, Vinay Kumar, Suresh Goyal, Akshita Bhat, Dr, Madhu Patni



PII: S1477-5131(17)30083-9

DOI: [10.1016/j.jpurol.2016.12.031](https://doi.org/10.1016/j.jpurol.2016.12.031)

Reference: JPUROL 2461

To appear in: *Journal of Pediatric Urology*

Received Date: 23 September 2016

Accepted Date: 23 December 2016

Please cite this article as: Bhat A, Bhat M, Kumar V, Goyal S, Bhat A, Patni M, The incidence of isolated penile torsion in north India: A study of 5,018 male neonates, *Journal of Pediatric Urology* (2017), doi: 10.1016/j.jpurol.2016.12.031.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The incidence of isolated penile torsion in north India: A study of 5,018 male neonates

Amilal Bhat ^{a,*}, Mahakshit Bhat ^b, Vinay Kumar ^a, Suresh Goyal ^c, Dr Akshita ^a, Madhu Patni ^d

^a Department of Urology, S.P. Medical College, Bikaner, Rajasthan, India

^b Department of Surgery, M.G. Medical College, Jaipur, Rajasthan India

^c Department of Urology, Dr S.N. Medical College Jodhpur, Rajasthan, India

^d Department of Obstetric and Gynecology, S.P. Medical College Bikaner, Rajasthan, India

* Corresponding author. Dr S.N. Medical College Jodhpur, C-15 sadul Ganj Bikaner, Jodhpur, Rajasthan 334003, India. Tel.: +91 151 222 6327; fax: +91 291 243 1987.

E-mail addresses: bhatamilal@gmail.com; amilalbhat@rediffmail.com

Summary

Objective

Congenital penile torsion is a three-dimensional deformity with helical rotation of the distal corporal bodies with the penile crurae remaining fixed to the pubic rami. The first case of congenital penile torsion (hypospadias) was described in 1857. Isolated penile torsion is an under-reported anomaly. The reported incidence of isolated penile torsion is 1.7–27% and severe torsion is 0.7%. There are no studies available from Indian subcontinent on the incidence of isolated penile torque. The objective of this study was to determine the overall incidence of isolated penile torque in a north Indian population.

Materials and methods

Download English Version:

<https://daneshyari.com/en/article/8811939>

Download Persian Version:

<https://daneshyari.com/article/8811939>

[Daneshyari.com](https://daneshyari.com)