


Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.pediatr-neonatology.com>

CASE REPORT

Fetal and Postnatal Magnetic Resonance Imaging of Unilateral Cystic Renal Dysplasia in a Neonate with Tuberous Sclerosis

Vineet Tyagi ^{a,*}, Eran Bornstein ^b, Robert Schacht ^c,
Shailee Lala ^d, Sarah Milla ^d^a *New York University School of Medicine, New York, NY, USA*^b *Department of Maternal-Fetal Medicine, Lenox Hill Hospital, New York, NY, USA*^c *Department of Pediatric Nephrology, New York University Langone Medical Center, New York, NY, USA*^d *Department of Radiology, New York University Langone Medical Center, New York, NY, USA*

Received Dec 29, 2012; received in revised form Sep 23, 2013; accepted Oct 22, 2013

Key Wordscystic renal disease;
tuberous sclerosis

Tuberous sclerosis (TS) is an autosomal dominant condition associated with mutations in the TSC1 and/or TSC2 genes. Clinical manifestations are multisystemic, and they often include lesions in the brain, skin, heart, kidneys, and bones. TSC2 gene mutations can be seen concomitantly with autosomal dominant polycystic kidney disease gene mutations. We present a case of a fetus with prenatal diagnosis of TS that had unique asymmetrical distribution of renal cystic disease. We describe the extensive work up with both fetal and neonatal magnetic resonance imaging with correlating images of the unilateral polycystic renal disease in addition to typical TS brain findings.

Copyright © 2014, Taiwan Pediatric Association. Published by Elsevier Taiwan LLC. All rights reserved.

1. Introduction

Tuberous sclerosis (TS) is a multisystemic disease that manifests with the growth of benign hamartomas in various

organ systems, including the central nervous system, kidneys, and skin. TS has an incidence rate of roughly 1 in 6000 newborns and affects over 1.5 million people globally. TS is an autosomal dominant disorder caused by the TSC1 and TSC2 genes, which are located on chromosomes 9 and 16, respectively. TS development generally requires a combination of events, the first of which is a germline mutation that results in a loss of heterozygosity for an allele of either gene. The subsequent events necessary for TS manifestation are somatic mutations that cause the loss of the other TS allele. Patients with TS often manifest brain lesions

Conflicts of interest: All contributing authors declare no conflicts of interest.

* Corresponding author. New York University School of Medicine, 545 First Avenue, 3H, New York, NY 10016, USA.

E-mail address: vineet.tyagi@med.nyu.edu (V. Tyagi).

1875-9572/\$36 Copyright © 2014, Taiwan Pediatric Association. Published by Elsevier Taiwan LLC. All rights reserved.

<http://dx.doi.org/10.1016/j.pedneo.2013.10.014>

Please cite this article in press as: Tyagi V, et al., Fetal and Postnatal Magnetic Resonance Imaging of Unilateral Cystic Renal Dysplasia in a Neonate with Tuberous Sclerosis, *Pediatrics and Neonatology* (2014), <http://dx.doi.org/10.1016/j.pedneo.2013.10.014>

(cortical/subcortical tubers, subependymal nodules, and subependymal giant cell tumors), angiomyolipomas, renal cysts, and skeletal lesions.¹

The diagnosis of TS requires a combination of either two major criteria, or one major and two minor criteria. TS may also be diagnosed based on the presence of TS-associated lesions in two different organs, or two differing types of lesions within the same organ. Probable TS is diagnosed by one major and one minor feature, and TS may be suspected in the presence of either one major or two minor features. The most common major features are cortical tubers, facial angiofibromas, nontraumatic ungual or periungual fibromas, hypomelanotic macules (>3), shagreen patches, subependymal nodules, subependymal giant cell tumors, cardiac rhabdomyoma (≥ 1), lymphangioleiomyomatosis, and renal angiomyolipomas. The minor features include pits in dental enamel, hamartomatous renal polyps, skeletal cysts, white matter migration tracts, gingival fibromas, nonrenal hamartoma, renal achromic patches, skin lesions, multiple renal cysts, and infantile spasms.²

2. Case report

A 28-year-old pregnant woman at 39 weeks of gestation was referred for fetal magnetic resonance imaging (MRI) to evaluate intracranial abnormalities that might support the prenatal diagnosis of TS. The fetus had just completed routine 3rd trimester sonographic evaluation where several fetal intracardiac masses with the sonographic appearance of rhabdomyomas were detected incidentally. Focused imaging of the fetal brain demonstrated subependymal nodules and cortical/subcortical signal abnormalities consistent with tubers (Figure 1). These findings confirmed

the diagnosis of TS. On scout images for the fetal MRI, an abnormal T2 hyperintensity was seen within the fetal abdomen. After dedicated imaging of the fetal brain, focused fetal body MRI images were obtained. It was noted that the right kidney was abnormally enlarged and had a lobulated mass-like appearance (Figure 2) with moderate T2/Half Fourier-Acquired Single Shot Turbo Spin Echo (HASTE) hyperintensity. No T1 hyperintensity was seen within the right kidney. The left kidney was normal in appearance. Given the unilaterality and the moderate T2 isointensity, a differential diagnosis for the appearance of the right kidney was given. Both solid and cystic masses such as mesoblastic nephroma and lipid poor angiomyolipoma were included. The couple denied any family history consistent with TS and the patient was counseled by both maternal fetal medicine and genetics specialists regarding these findings.

Following uncomplicated spontaneous vaginal delivery, postnatal MRI of the brain confirmed the prenatal findings of subependymal nodules and cortical/subcortical tubers (Figure 1). Postnatal sonography of the kidneys demonstrated an enlarged right kidney with a polycystic appearance and a left kidney that appeared to be normal (Figure 2). The neonate was discharged home in good medical condition and follow-up was scheduled in a specialized TS clinic.

A follow-up ultrasound was performed at age 2 months, and there was an overall decrease in the size of the multiple cysts; however, there was a persistent unilateral polycystic appearance (Figure 3). MRI of the kidneys demonstrated a unilateral segmentally distributed polycystic appearance to the right kidney interspersed with more normally appearing renal parenchyma with a preserved corticomedullary appearance in focal areas. The right




Figure 1 (A,B) Axial Half Fourier-Acquired Single Shot Turbo Spin Echo (HASTE) and (C,D) Axial T1 images from a fetal magnetic resonance scan demonstrating T2 hypointense and T1 hyperintense subependymal nodules (white arrow with black outline) and cortical/subcortical tubers (white arrows). (E,F) Axial T2 and (G,H) Axial T1 images from the postnatal magnetic resonance imaging demonstrating T2 hypointense and T1 hyperintense subependymal nodules (white arrow with black outline) and cortical/subcortical tubers (white arrows).

Download English Version:

<https://daneshyari.com/en/article/8813578>

Download Persian Version:

<https://daneshyari.com/article/8813578>

[Daneshyari.com](https://daneshyari.com)