

## Accepted Manuscript

Bipolar features in major depressive disorder: Results from the Iranian mental health survey (IranMHS)

Masoumeh Amin-Esmaeili , Abbas Motevalian ,  
Afarin Rahimi-Movaghar , Ahmad Hajebi , Vandad Sharifi ,  
Ramin Mojtabai , Shahrokh S Gudarzi

PII: S0165-0327(18)30954-6  
DOI: <https://doi.org/10.1016/j.jad.2018.08.014>  
Reference: JAD 10011



To appear in: *Journal of Affective Disorders*

Received date: 4 May 2018  
Revised date: 11 July 2018  
Accepted date: 7 August 2018

Please cite this article as: Masoumeh Amin-Esmaeili , Abbas Motevalian , Afarin Rahimi-Movaghar , Ahmad Hajebi , Vandad Sharifi , Ramin Mojtabai , Shahrokh S Gudarzi , Bipolar features in major depressive disorder: Results from the Iranian mental health survey (IranMHS), *Journal of Affective Disorders* (2018), doi: <https://doi.org/10.1016/j.jad.2018.08.014>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Highlights

- More than one in five of the individuals with MDD experience bipolar features.
- The bipolar features are associated with a distinct demographic and clinical profile.
- Anxiety disorders were independently associated with the bipolar features.
- History of psychotic symptoms was also independently associated with the features.

Download English Version:

<https://daneshyari.com/en/article/8814999>

Download Persian Version:

<https://daneshyari.com/article/8814999>

[Daneshyari.com](https://daneshyari.com)