

## Accepted Manuscript

Persistent Depressive and Grief Symptoms for up to 10 years Following Perinatal Loss: Involvement of Negative Cognitions

Kossigan Kokou-Kpolou , Olga Megalakaki , Nicolas Nieuviarts

PII: S0165-0327(18)30814-0  
DOI: <https://doi.org/10.1016/j.jad.2018.08.063>  
Reference: JAD 10061



To appear in: *Journal of Affective Disorders*

Received date: 16 April 2018  
Revised date: 8 June 2018  
Accepted date: 12 August 2018

Please cite this article as: Kossigan Kokou-Kpolou , Olga Megalakaki , Nicolas Nieuviarts , Persistent Depressive and Grief Symptoms for up to 10 years Following Perinatal Loss: Involvement of Negative Cognitions, *Journal of Affective Disorders* (2018), doi: <https://doi.org/10.1016/j.jad.2018.08.063>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Highlights

- For the first time, cognitive mechanisms underlying the perinatal grief process were examined.
- Maladaptive cognitions about life, the future and the world increased perinatal grief symptoms.
- Specific maladaptive cognition about the world increased depressive symptoms following perinatal loss.
- Maladaptive cognitions varied depending on the type of loss.

Download English Version:

<https://daneshyari.com/en/article/8815008>

Download Persian Version:

<https://daneshyari.com/article/8815008>

[Daneshyari.com](https://daneshyari.com)