

Accepted Manuscript

The relations between empathy, guilt, shame and depression in inpatient adolescents

Malgorzata Gambin , Carla Sharp

PII: S0165-0327(17)32536-3
DOI: <https://doi.org/10.1016/j.jad.2018.08.068>
Reference: JAD 10066



To appear in: *Journal of Affective Disorders*

Received date: 9 December 2017
Revised date: 21 June 2018
Accepted date: 12 August 2018

Please cite this article as: Malgorzata Gambin , Carla Sharp , The relations between empathy, guilt, shame and depression in inpatient adolescents, *Journal of Affective Disorders* (2018), doi: <https://doi.org/10.1016/j.jad.2018.08.068>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Generalized shame, contextual shame and generalized guilt mediate the relationship between affective empathy and depressive symptoms in inpatient adolescents.
- Cognitive empathy is associated the most strongly with contextual guilt and not related to depressive symptoms.
- Shame and generalized guilt associated with affective sharing should be considered as possible targets for therapeutic and preventive interventions for adolescents with high levels of depressive symptoms.

Download English Version:

<https://daneshyari.com/en/article/8815015>

Download Persian Version:

<https://daneshyari.com/article/8815015>

[Daneshyari.com](https://daneshyari.com)