

Accepted Manuscript

Treating a broader range of depressed adolescents with combined therapy

Simon Foster PhD , Prof. Meichun Mohler-Kuo ScD

PII: S0165-0327(18)30384-7
DOI: <https://doi.org/10.1016/j.jad.2018.08.027>
Reference: JAD 10024



To appear in: *Journal of Affective Disorders*

Received date: 23 February 2018
Revised date: 26 July 2018
Accepted date: 7 August 2018

Please cite this article as: Simon Foster PhD , Prof. Meichun Mohler-Kuo ScD , Treating a broader range of depressed adolescents with combined therapy, *Journal of Affective Disorders* (2018), doi: <https://doi.org/10.1016/j.jad.2018.08.027>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- **Response distributions of depressed adolescents were compared across treatments**
- **Cognitive-behavioral therapy was no more effective than placebo**
- **Fluoxetine was effective only in the middle range of response levels**
- **Combining cognitive-behavioral therapy and fluoxetine was broadly effective**
- **Combining the two treatments was also effective in low responders**

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8815023>

Download Persian Version:

<https://daneshyari.com/article/8815023>

[Daneshyari.com](https://daneshyari.com)