## **Accepted Manuscript**

Availability and Use of Mental Health Services in European Countries: Influence on National Suicide Rates


Daniel König, Matthäus Fellinger, Nathalie Pruckner, Barbara Hinterbuchinger, Georg Dorffner, Andreas Gleiss, Sandra Vyssoki, Benjamin Vyssoki

PII: S0165-0327(18)30586-X DOI: 10.1016/j.jad.2018.06.042

Reference: JAD 9900

To appear in: Journal of Affective Disorders

Received date: 9 April 2018
Revised date: 13 June 2018
Accepted date: 15 June 2018



Please cite this article as: Daniel König, Matthäus Fellinger, Nathalie Pruckner, Barbara Hinterbuchinger, Georg Dorffner, Andreas Gleiss, Sandra Vyssoki, Benjamin Vyssoki, Availability and Use of Mental Health Services in European Countries: Influence on National Suicide Rates, *Journal of Affective Disorders* (2018), doi: 10.1016/j.jad.2018.06.042

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

### ACCEPTED MANUSCRIPT

## Highlights

- No statistically significant influence of mental health care variables on SMR remained after correcting for country, time and economic variables
- Available data predicted less than 10% of the variability of the suicide mortality rate.
- Further research is needed to implement preventive measures for individuals at risk of suicide.



#### Download English Version:

# https://daneshyari.com/en/article/8815075

Download Persian Version:

https://daneshyari.com/article/8815075

<u>Daneshyari.com</u>