

Accepted Manuscript

Comparative risk of lipophilic and hydrophilic statins on incident depression: A retrospective cohort study

Chintan V. Dave , Almut G. Winterstein , Haesuk Park ,
Robert L. Cook , Abraham G. Hartzema

PII: S0165-0327(18)30810-3
DOI: [10.1016/j.jad.2018.06.021](https://doi.org/10.1016/j.jad.2018.06.021)
Reference: JAD 9867



To appear in: *Journal of Affective Disorders*

Received date: 16 April 2018
Revised date: 21 May 2018
Accepted date: 4 June 2018

Please cite this article as: Chintan V. Dave , Almut G. Winterstein , Haesuk Park , Robert L. Cook , Abraham G. Hartzema , Comparative risk of lipophilic and hydrophilic statins on incident depression: A retrospective cohort study, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.06.021](https://doi.org/10.1016/j.jad.2018.06.021)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Lipophilic statins cross the blood-brain barrier and associated with mood disorders
- 1:1 propensity score matched cohort of 299,298 patients initiating a lipophilic v hydrophilic statin was created
- The study did not find an increase in the risk of incident depression in patients initiating a lipophilic statin

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8815340>

Download Persian Version:

<https://daneshyari.com/article/8815340>

[Daneshyari.com](https://daneshyari.com)