


Accepted Manuscript

Classification of Attempted Suicide by Cluster Analysis: A Study of 888 Suicide Attempters Presenting to the Emergency Department

Hyeyoung Kim , Bora Kim , Se Hyun Kim , C. Hyung Keun Park , Eun Young Kim , Yong Min Ahn

PII: S0165-0327(17)31946-8
DOI: [10.1016/j.jad.2018.04.001](https://doi.org/10.1016/j.jad.2018.04.001)
Reference: JAD 9655



To appear in: *Journal of Affective Disorders*

Received date: 29 September 2017
Revised date: 18 February 2018
Accepted date: 2 April 2018

Please cite this article as: Hyeyoung Kim , Bora Kim , Se Hyun Kim , C. Hyung Keun Park , Eun Young Kim , Yong Min Ahn , Classification of Attempted Suicide by Cluster Analysis: A Study of 888 Suicide Attempters Presenting to the Emergency Department, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.04.001](https://doi.org/10.1016/j.jad.2018.04.001)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Cluster analysis extracted two distinct subgroups of Korean suicide attempters.
- Eighty-five percent of suicide attempts occurred impulsively.
- The remaining 14.6% of attempts were more lethal and well-planned.
- The two clusters showed differences in several demographic and clinical variables.

Download English Version:

<https://daneshyari.com/en/article/8815349>

Download Persian Version:

<https://daneshyari.com/article/8815349>

[Daneshyari.com](https://daneshyari.com)