


Accepted Manuscript

Prior suicide attempts predict worse clinical and functional outcomes in young people attending a mental health service

Frank Iorfino , Daniel. F. Hermens , Shane Cross ,
Natalia Zmicerevska , Alissa Nichles , Josine Groot ,
Adam Guastella , Elizabeth Scott , Ian. B. Hickie

PII: S0165-0327(17)32606-X
DOI: [10.1016/j.jad.2018.06.032](https://doi.org/10.1016/j.jad.2018.06.032)
Reference: JAD 9889



To appear in: *Journal of Affective Disorders*

Received date: 4 March 2018
Revised date: 7 May 2018
Accepted date: 12 June 2018

Please cite this article as: Frank Iorfino , Daniel. F. Hermens , Shane Cross , Natalia Zmicerevska , Alissa Nichles , Josine Groot , Adam Guastella , Elizabeth Scott , Ian. B. Hickie , Prior suicide attempts predict worse clinical and functional outcomes in young people attending a mental health service, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.06.032](https://doi.org/10.1016/j.jad.2018.06.032)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

HIGHLIGHTS

- Prior suicide attempts indicate greater vulnerability to worse clinical and function outcomes.
- Prior suicide attempts were associated with the onset of bipolar disorder.
- New-onset suicide attempts were more common than recurrent attempts during follow-up.
- Recurrent suicide attempts were associated with being older and an alcohol/substance use disorder.

Download English Version:

<https://daneshyari.com/en/article/8815352>

Download Persian Version:

<https://daneshyari.com/article/8815352>

[Daneshyari.com](https://daneshyari.com)