


Accepted Manuscript

Task-positive and task-negative networks in major depressive disorder: A combined fMRI and EEG study

Gennady G. Knyazev , Alexander N. Savostyanov ,
Andrey V. Bocharov , Ivan V. Brak , Evgeny A. Osipov ,
Elena A. Filimonova , Alexander E. Saprigyn , Lyubomir I. Aftanas

PII: S0165-0327(17)32134-1
DOI: [10.1016/j.jad.2018.04.003](https://doi.org/10.1016/j.jad.2018.04.003)
Reference: JAD 9657



To appear in: *Journal of Affective Disorders*

Received date: 14 October 2017
Revised date: 23 February 2018
Accepted date: 2 April 2018

Please cite this article as: Gennady G. Knyazev , Alexander N. Savostyanov , Andrey V. Bocharov , Ivan V. Brak , Evgeny A. Osipov , Elena A. Filimonova , Alexander E. Saprigyn , Lyubomir I. Aftanas , Task-positive and task-negative networks in major depressive disorder: A combined fMRI and EEG study, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.04.003](https://doi.org/10.1016/j.jad.2018.04.003)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Depression-related dominance of DMN over TPN was tested using fMRI and EEG data.
- The dominance was observed in major emotion and attention regulation centers.
- These centers are more ready to respond to self-related thoughts than to environment.

Download English Version:

<https://daneshyari.com/en/article/8815357>

Download Persian Version:

<https://daneshyari.com/article/8815357>

[Daneshyari.com](https://daneshyari.com)