

Accepted Manuscript

Improvement of postpartum depression and psychosis after electroconvulsive therapy: – a population-based study with a matched comparison group

Sara Rundgren , Ole Brus , Ullvi Båve , Mikael Landen ,
Johan Lundberg , Pia Nordanskog , Axel Nordenskjöld

PII: S0165-0327(18)30206-4
DOI: [10.1016/j.jad.2018.04.043](https://doi.org/10.1016/j.jad.2018.04.043)
Reference: JAD 9697



To appear in: *Journal of Affective Disorders*

Received date: 2 February 2018
Revised date: 29 March 2018
Accepted date: 4 April 2018

Please cite this article as: Sara Rundgren , Ole Brus , Ullvi Båve , Mikael Landen , Johan Lundberg , Pia Nordanskog , Axel Nordenskjöld , Improvement of postpartum depression and psychosis after electroconvulsive therapy: – a population-based study with a matched comparison group, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.04.043](https://doi.org/10.1016/j.jad.2018.04.043)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- The response rate to ECT is higher in the postpartum period than outside this period.
- Severe forms of postpartum depression and psychosis respond well to ECT.
- ECT is a valid treatment option for psychosis and severe depression postpartum.

Download English Version:

<https://daneshyari.com/en/article/8815373>

Download Persian Version:

<https://daneshyari.com/article/8815373>

[Daneshyari.com](https://daneshyari.com)