

Accepted Manuscript

Associations between cognitive functioning, mood symptoms and coping styles in older age bipolar disorder

Nadine P.G. Paans , Annemiek Dols , Hannie C. Comijs ,
Max L. Stek , Sigfried N.T.M. Schouws

PII: S0165-0327(18)30192-7
DOI: [10.1016/j.jad.2018.04.052](https://doi.org/10.1016/j.jad.2018.04.052)
Reference: JAD 9706



To appear in: *Journal of Affective Disorders*

Received date: 29 January 2018
Revised date: 29 March 2018
Accepted date: 4 April 2018

Please cite this article as: Nadine P.G. Paans , Annemiek Dols , Hannie C. Comijs , Max L. Stek , Sigfried N.T.M. Schouws , Associations between cognitive functioning, mood symptoms and coping styles in older age bipolar disorder, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.04.052](https://doi.org/10.1016/j.jad.2018.04.052)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Better executive functioning is associated to more active coping.
- Less depressive symptoms are associated to more active coping.
- Mood is a more important contributor to active coping as compare to cognition.
- No associations between cognition, mood and passive coping were found.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8815416>

Download Persian Version:

<https://daneshyari.com/article/8815416>

[Daneshyari.com](https://daneshyari.com)