Accepted Manuscript



Looking for bipolarity in antidepressant discontinuation manic states: update and diagnostic considerations of the phenomenon

Sandra Abou Kassm , Wadih Naja

PII: S0165-0327(18)30319-7
DOI: 10.1016/j.jad.2018.04.080
Reference: JAD 9735

To appear in: *Journal of Affective Disorders*

Received date: 14 February 2018
Revised date: 8 April 2018
Accepted date: 14 April 2018

Please cite this article as: Sandra Abou Kassm , Wadih Naja , Looking for bipolarity in antidepressant discontinuation manic states: update and diagnostic considerations of the phenomenon, *Journal of Affective Disorders* (2018), doi: 10.1016/j.jad.2018.04.080

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Highlights

- Almost any antidepressant can cause a manic state upon its discontinuation
- Discontinuation mania may involve Catecholamines, Acetylcholine and Serotonin
- More than half of patients with discontinuation mania can be included in the bipolar spectrum
- Vulnerability to discontinuation mania can be referred to as bipolar III 1/4

ACCEPTED MANUSCRIPT

Download English Version:

https://daneshyari.com/en/article/8815472

Download Persian Version:

https://daneshyari.com/article/8815472

Daneshyari.com