

Accepted Manuscript

Work intensity in men and work-related emotional demands in women are associated with increased suicidality among persons attending primary care

N Younès , M Rivière , L Plancke , A Leroyer , T Blanchon ,
M Azevedo Da Silva , M Melchior

PII: S0165-0327(17)32681-2
DOI: [10.1016/j.jad.2018.04.075](https://doi.org/10.1016/j.jad.2018.04.075)
Reference: JAD 9730



To appear in: *Journal of Affective Disorders*

Received date: 22 December 2017
Revised date: 6 April 2018
Accepted date: 10 April 2018

Please cite this article as: N Younès , M Rivière , L Plancke , A Leroyer , T Blanchon , M Azevedo Da Silva , M Melchior , Work intensity in men and work-related emotional demands in women are associated with increased suicidality among persons attending primary care, *Journal of Affective Disorders* (2018), doi: [10.1016/j.jad.2018.04.075](https://doi.org/10.1016/j.jad.2018.04.075)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Among working individuals consulting a primary care physician, 8.0% reported suicidal ideations in the preceding month.
- Suicidality is significantly associated with work intensity in men and with work-related emotional demands in women.
- To prevent suicide, a global strategy should include GPs.

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/8815476>

Download Persian Version:

<https://daneshyari.com/article/8815476>

[Daneshyari.com](https://daneshyari.com)