


Author's Accepted Manuscript

Predictors of hospitalization length of stay among re-admitted treatment-resistant bipolar disorder inpatients

M. Fornaro, F. Iasevoli, S. Novello, A. Fusco, A. Anastasia, D. De Berardis, A. Valchera, A. de Bartolomeis



PII: S0165-0327(17)32048-7
DOI: <https://doi.org/10.1016/j.jad.2017.12.009>
Reference: JAD9417

To appear in: *Journal of Affective Disorders*

Received date: 2 October 2017
Revised date: 8 November 2017
Accepted date: 5 December 2017

Cite this article as: M. Fornaro, F. Iasevoli, S. Novello, A. Fusco, A. Anastasia, D. De Berardis, A. Valchera and A. de Bartolomeis, Predictors of hospitalization length of stay among re-admitted treatment-resistant bipolar disorder inpatients, *Journal of Affective Disorders*, <https://doi.org/10.1016/j.jad.2017.12.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Predictors of hospitalization length of stay among re-admitted treatment-resistant bipolar disorder inpatients[★]

M. Fornaro^{*}, F. Iasevoli, S. Novello, A. Fusco, A. Anastasia, D. De Berardis, A.

Valchera, A. de Bartolomeis

dott.fornaro@gmail.com

felix_ias@hotmail.com

Stefano.novello1@gmail.com

andreausco1990@gmail.com

annalisa.anastasia@yahoo.it

domenico.deberardis@asliteramo.it

a.valchera@ospedaliere.it

adebarto@unina.it

^{*}Correspondence to: Dr. Michele Fornaro, MD, PhD, Section of Psychiatry - Unit on Treatment Resistant Disorders, Department of Neuroscience, Reproductive Sciences and Odontostomatology University School of Medicine Federico II, Naples, Via Pansini 5, 80131 Napoli Italy. Tel.: +081 7463070. dott.fornaro@gmail.com

Abstract

Background

Hospitalization accounts for significant health care resource utilization for treatment-resistant Bipolar Disorder (BD), especially among frequent users of acute inpatient

[★] “*Hospitalization length among TRBD inpatients*”

Download English Version:

<https://daneshyari.com/en/article/8815801>

Download Persian Version:

<https://daneshyari.com/article/8815801>

[Daneshyari.com](https://daneshyari.com)