

Author's Accepted Manuscript

Alcohol-induced disinhibition is associated with impulsivity, depression, and suicide attempt: a nationwide community sample of Korean adults

Kwan Woo Choi, Eun Jin Na, Jin Pyo Hong, Maeng Je Cho, Maurizio Fava, David Mischoulon, Hana Cho, Hong Jin Jeon



PII: S0165-0327(17)31313-7
DOI: <https://doi.org/10.1016/j.jad.2017.11.001>
Reference: JAD9315

To appear in: *Journal of Affective Disorders*

Received date: 30 June 2017
Revised date: 31 August 2017
Accepted date: 2 November 2017

Cite this article as: Kwan Woo Choi, Eun Jin Na, Jin Pyo Hong, Maeng Je Cho, Maurizio Fava, David Mischoulon, Hana Cho and Hong Jin Jeon, Alcohol-induced disinhibition is associated with impulsivity, depression, and suicide attempt: a nationwide community sample of Korean adults, *Journal of Affective Disorders*, <https://doi.org/10.1016/j.jad.2017.11.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Alcohol-induced disinhibition is associated with impulsivity, depression,
and suicide attempt: a nationwide community sample of Korean adults[☆]**

Kwan Woo Choi, MD^a, Eun Jin Na, MD^{a,b}, Jin Pyo Hong, MD, Ph.D^a, Maeng Je Cho, MD, Ph.D^c, Maurizio Fava, MD^d, David Mischoulon, MD, Ph.D^d, Hana Cho, MD, Ph.D^e, Hong Jin Jeon, MD, Ph.D^{a,d,f*}

^aDepartment of Psychiatry, Depression Center, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

^bKorean Psychological Autopsy Center (KPAC), Seoul, Korea

^cDepartment of Psychiatry, Seoul National University, College of Medicine, Seoul, Korea

^dDepression Clinical and Research Program, Massachusetts General Hospital, Harvard Medical School, Boston, USA

^eDepartment of Physiology, Sungkyunkwan University School of Medicine, Samsung Biomedical Research Institute, Suwon 440-746, Korea

^fDepartment of Health Sciences & Technology, Department of Medical Device Management & Research, and Department of Clinical Research Design & Evaluation, Samsung Advanced Institute for Health Sciences & Technology (SAIHST), Sungkyunkwan University, Seoul, Korea

*Corresponding author at: Department of Psychiatry, Depression Center, Samsung Medical Center, Sungkyunkwan University School of Medicine, #81 Irwon-ro, Gangnam-gu, Seoul, South Korea, 06351. Tel.: +82 2 3410 3586; fax: +82 2 3410 0050. jeonhj@skku.edu

Abstract

Background

Alcohol-induced disinhibition (AID) is defined as a “loss of restraint over some form of behavior after drinking alcohol regardless of its amount”. Although many previous studies

[☆] All authors agreed to this submission.

The material is original research, has not been previously published and has not been submitted for publication elsewhere while under consideration.

Download English Version:

<https://daneshyari.com/en/article/8815893>

Download Persian Version:

<https://daneshyari.com/article/8815893>

[Daneshyari.com](https://daneshyari.com)