

Letter to the Editor



Clustering of Non-communicable Diseases Risk Factors in Healthy Adults Aged 35 Years and Older in Shenzhen, China*

NI Wen Qing¹, XU Jian¹, LIU Min², LIU Xiao Li¹, YANG Li Chen³, ZHUO Zhi Peng¹,
YUAN Xue Li¹, SONG Jin Ping¹, CHI Hong Shan¹, and BAI Ya Min^{2,#}

We assessed the prevalence of non-communicable diseases (NCDs) risk factors with a focus on their clustering among healthy adults in Shenzhen, China. Data from the 2011 China Health and Nutrition Survey, comprising a regionally representative sample of 806 healthy adults aged 35 years or older, were obtained to determine the prevalence of five risk factors for NCDs. The prevalence of current smoking, central obesity, impaired fasting glucose, borderline hypertension, and borderline high total cholesterol was 19.97%, 28.29%, 4.47%, 10.55%, and 36.10%, respectively. A total 63.77% of participants had at least one risk factor. Upon examination of risk factor clustering, we observed that 7.57% of participants had at least three risk factors. Using this threshold as a cutoff, clustering of risk factors was associated with sex [odds ratio (OR) = 3.336, 95% confidence interval (CI): 1.782 to 6.246], physical activity (OR = 1.913, 95% CI: 1.009 to 3.628), and BMI (OR = 7.376, 95% CI: 3.812 to 14.270). The prevalence of risk factors for NCDs is fairly high among healthy adults in Shenzhen, with a clustering tendency.

Key words: Non-communicable diseases; Healthy adults; Prevalence; Risk factors

The leading causes of death among people in China are non-communicable diseases (NCDs)^[1]. Tobacco smoking, central obesity, borderline high total cholesterol (TC), borderline hypertension, and raised fasting glucose levels are well known shared modifiable risk factors for the major NCDs. The national standards for the prevention and control of

chronic diseases, issued by the Ministry of Health of the People's Republic of China, lists these five risk factors as critical to the control of NCDs. These risk factors are more prevalent and easier to detect than NCDs. Therefore, assessing epidemiological status by identifying the distribution of risk factors among different population in a region is the first of three planning steps recommended by the World Health Organization (WHO) for the prevention and control of NCDs and their risk factors. Information on a regional risk factor profile could also aid in predicting the future burden of disease. This in turn helps to make a strong case for high-level advocacy and constitutes an evidence base for planning interventions at policy, environmental, and health system levels.

There have been reports from the United States on clustering of risk factors among the population^[2]. This clustering phenomenon may predispose Americans to a higher burden of NCDs than populations with lower clustering tendencies. At the same time, previous studies have highlighted the need for local, high-quality epidemiological data on the burden of NCDs and their risk factors, particularly in Shenzhen where such data are scarce^[3]. However, systematic analysis of the clustering of all major risk factors is lacking, making it difficult to establish a regional representation of healthy people in Shenzhen. Therefore, we conducted a cross-sectional study to obtain a comprehensive profile of the epidemiologic distribution of selected risk factors for NCDs among healthy residents of Shenzhen.

doi: 10.3967/bes2017.087

*This study was supported by National Project of NCDs High-risk Population Health Management, Center for Chronic and Non-communicable Diseases Control and Prevention, ChinaCDC (Grant No. 2013085); The Science and Technology Planning Project of Shenzhen City, Guangdong Province, China (Grant No. 201602005).

1. Shenzhen Center for Chronic Disease Control, Shenzhen 518020, Guangdong, China; 2. Center for Chronic and Non-communicable Diseases Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 100050, China; 3. Key laboratory of Trace Element Nutrition of the Ministry of Health, National Institute of Nutrition for Health, Chinese Center for Disease Control and Prevention, Beijing 100050, China

The present cross-sectional study was conducted as a part of the China Health and Nutrition Survey at the Center for Chronic Disease Control of Shenzhen. The ethics committee of the Center for Chronic Disease Control of Shenzhen approved the study. The survey is described in detail elsewhere^[4]. In brief, the survey used a multistage stratified cluster random sampling method to select a representative sample of permanent residents in eight different districts of the city of Shenzhen. The first stage of sampling was stratified by district and population distribution on the basis of Shenzhen population data from 2010, which involved the random selection of 12 streets from the 8 districts. In the second stage, one residential community was randomly selected from each selected street. In the third stage, about 75 households were randomly selected from the residential communities selected in the earlier stage. In the last stage, eligible family member from each designated household was recruited.

The eligibility criteria were set as follows: a man or non-pregnant woman; aged 35 years or above; living in Shenzhen for more than 5 years; and confirmed not to have hypertension, diabetes, high TC, cancer, severe liver disease, systemic lupus erythematosus, or kidney diseases. From February to July 2011, according to the eligibility criteria, a total of 882 adults were selected from designated households and invited to participate in the study. We excluded 76 individuals because they did not provide questionnaire-derived information or fasting blood or anthropometry data. Finally, 806 participants were enrolled in the study. The population size was relatively small because of the limited number of healthy adults in the region. Structured questionnaires were used to collect information on sociodemographic characteristic variables and health parameters. Height, body weight, waist circumference (WC), and blood pressure of all participants were recorded. Fasting blood glucose (FBG) and TC concentrations were measured using commercially available kits.

Each participant was also categorized as a habitual drinker, non-habitual drinker, or non-drinker^[4]. In this study, the term 'moderate- to vigorous-intensity physical activity' was defined as physical activity causing at least some sweating or shortness of breath, whereas the term 'light physical activity' was defined as causing no sweating or shortness of breath^[4]. Physical activity was further categorized as moderate- to vigorous-intensity

physical activity performed once a week or more. Current smoking was defined as self-reported use of tobacco products every day or on some days, at the time the survey was conducted^[4]. We defined impaired fasting glucose as a value of 6.1-6.9 mmol/L^[5], and borderline hypertension was defined as systolic blood pressure (SBP) 130-139 mmHg or diastolic blood pressure (DBP) 85-89 mmHg^[6]. Borderline high TC was defined as having a TC value ≥ 5.2 mmol/L and < 6.2 mmol/L. Body mass index (BMI) was defined as weight (kg) divided by the square of height (m) and was classified into the following categories according to the accepted Chinese BMI standard: underweight (BMI < 18.5 kg/m²), normal weight ($18.5 \leq$ BMI < 24.0 kg/m²), overweight ($24.0 \leq$ BMI < 28.0 kg/m²), and obese (BMI ≥ 28.0 kg/m²). Two sets of cutoff values were used to define central obesity in terms of WC: ≥ 90 cm for men and ≥ 85 cm for women on the basis of specific guidelines for central obesity.

The mean and standard deviation of continuous variables were expressed as mean \pm SD, and percentage was calculated for the categorical variables. Age and other anthropometric characteristic differences between groups by sex were determined using a Student *t*-test. The chi-squared test or Fisher's exact test was used to determine significant differences in proportions among categorical variables. We then determined the number of selected risk factors for each participant at the time of the survey (from 0 to 5). In addition, Spearman analysis was performed to identify possible predictors of the number of selected risk factors. Multiple logistic regression analysis was done to obtain odds ratios (ORs) and their confidence intervals (CIs) for clustering of three or more selected risk factors. Statistical analysis of the data was performed using IBM SPSS version 21.0 software (IBM Corp., Armonk, NY, USA). All statistical analyses were two-tailed and $P < 0.05$ was considered statistically significant.

Table 1 summarizes the sociodemographic and other characteristics of the 806 participants who were included in the final analysis. Of the included participants, 44.17% were men, 55.83% were women, the mean age was 48.94 ± 11.32 years, 51.24% had attained a senior high school or higher education, 94.54% were currently married, and 48.26% were employed. The rate of current smoking was 19.98%, habitual drinking was 6.70%, and 83.87% of participants reported engaging in regular physical activity. In terms of anthropometric

Download English Version:

<https://daneshyari.com/en/article/8817530>

Download Persian Version:

<https://daneshyari.com/article/8817530>

[Daneshyari.com](https://daneshyari.com)