



## Case report

# The role and safety of endobronchial ultrasound-guided transbronchial needle aspiration in the diagnosis and management of infected bronchogenic mediastinal cysts in adults

Hazim S. Bukamur<sup>a,\*</sup>, Emad Alkhankan<sup>a</sup>, Haitem M. Mezughi<sup>b</sup>, Nancy J. Munn<sup>a</sup>,  
Yousef R. Shweihat<sup>a</sup>

<sup>a</sup> Department of Pulmonary and Critical Care Medicine, Marshall University, Joan C. Edwards School of Medicine and the Huntington Veterans Administration Medical Center, Huntington, WV 25701, USA

<sup>b</sup> Department of Internal Medicine, Marshall University, Joan C. Edwards School of Medicine, Huntington, WV 25701, USA

## ARTICLE INFO

## Keywords:

Endobronchial ultrasound  
Fine needle aspiration  
Bronchogenic cyst

## ABSTRACT

Bronchogenic and other duplication cysts are congenital abnormalities that can present at any age including adulthood years. They are usually asymptomatic and discovered incidentally on radiological imaging of the chest. They are commonly treated by surgical resection. Recently, endobronchial ultrasound has been used to assist in diagnosis when radiologic imaging is not definitive. Endobronchial ultrasound has been used rarely to drain infected cysts, a rare complication of the bronchogenic cyst. We present a unique case of an infected large bronchogenic cyst treated with endobronchial ultrasound drainage combined with conservative medical therapy. We also review the scarce available literature describing such an approach and its potential complications and add recommendations based on our experience in managing these anomalies.

## 1. Introduction

Bronchogenic cysts represent a spectrum of bronchopulmonary malformations that result from an abnormal budding of the tracheobronchial tree during fetal development. Most bronchogenic cysts (BCs) are located in the mediastinum with 15%–20% occurring in the lung parenchyma, mostly in lower lobes. Communication with the airway is rare but can predispose to cyst infection [1–4]. Most patients with mediastinal BC are asymptomatic and found incidentally on radiological imaging of the chest. The most frequent symptoms are cough (the most common presentation), dyspnea, fever and pain [1,3,4]. Some patients present with complications, including respiratory distress from airway compression, infection, vascular compression, malignant transformation or airway fistulae [3,4]. Mediastinal BC is usually diagnosed with CT scan or MRI revealing fluid density or resonance in the abnormality detected [4]. BCs are treated with resection even when asymptomatic due to the small increased risk of malignancy [5]. However, not all authors agree with this invasive approach [6]. Surgical drainage and resection of the cyst has been the classic approach for complicated or large mediastinal cysts [7]. This is reported to prevent rupture and seepage of the fluids, especially if infected, into the mediastinum and pleural space [7]. Moreover, minimally invasive surgical


methods, such as mediastinoscopy, percutaneous drainage approach and endobronchial ultrasound-transbronchial needle aspiration (EBUS-TBNA) now appear safe [8–11].

## 2. Case presentation

A 33-year-old male had symptoms of cough, sputum production and shortness of breath 4 months before admission. He presented with symptoms of upper respiratory tract infection with fever, chills, productive cough and retrosternal chest pain. He had these symptoms start shortly after a previous similar episode. He denied weight loss. On exam, the patient was tachycardic, with a heart rate of 150 bpm. Otherwise his exam was normal. Laboratory workup revealed leukocytosis and normal chemistry panel and thyroid function panel. Blood cultures were negative. EKG showed atrial fibrillation with rapid ventricular response that converted to sinus rhythm with medical treatment. His CXR showed a widened mediastinum and possible lung infiltrates. CT of the chest revealed a mediastinal homogeneous mass with fluid density, located within the prevascular space and sub-carinal area measuring approximately  $8.6 \times 2.4$  cm (Fig. 1-a, b, c). It was thought his atrial fibrillation was directly related to compression of the cyst on the atria or related to the inflammation/irritation caused by the

\* Corresponding author.

E-mail address: [hazimbukamur@gmail.com](mailto:hazimbukamur@gmail.com) (H.S. Bukamur).



**Fig. 1.** A and B showing mid and lower sections through the fluid filled bronchogenic cyst in the subcarinal location. C shows the uppermost border of the cyst and its extension into the pre-vascular space (Arrow).

infected cyst on the atrial wall. We performed EBUS flexible bronchoscopy in the operating room under general anesthesia. A round hypoechoic structure, suspicious for a septated cystic lesion was detected. A fine needle aspiration was performed under direct visualization removing 50 ml of thick yellow material after single puncture of the cyst. Cytology from the aspirate was negative; culture grew *Streptococcus viridans*, *Coryneform bacteria* and *Prevotella melaninogenica*. The patient was treated with antibiotics. He improved significantly and was discharged to be evaluated for resection after infection was controlled. The patient was evaluated by a multidisciplinary approach and referred to an advanced thoracic surgery quaternary center. Due to the complexity and location of the cyst and resolution of symptoms at time of the evaluation, the conservative approach of follow up was elected. He was treated with 4 weeks of Amoxicillin/clavulanic acid and was followed up clinically at 3 months intervals. He had CT scans every 6 months for the first year and yearly thereafter.

### 3. Discussion

We believe the cyst we encountered in this case to be a bronchogenic cyst given the location in the subcarinal and prevascular mediastinal location. The absence of other systemic symptoms or manifestations make acquired cystic diseases like lymphoma or hydatid cyst unlikely. Esophageal cysts are usually tubular and in close association with the esophagus unlike our case that is extensive and extends to the prevascular space. Cytologic examination of aspirate did not reveal any lymphocytic or thymic tissue effectively ruling out lymphomas and thymus related cysts. Other etiologies like pericardial cyst is highly unlikely based only on location.

Infection is a rare complication of bronchogenic cysts but it is among the most serious, potentially life-threatening conditions. Early presentation of cyst infection is vague and nonspecific. It often will progress insidiously over time [1] as probably occurred in our patient. Delayed treatment increases risk of complications. Thus, early diagnosis and prompt management would likely assure a good prognosis [1,12].

There is little argument on the usefulness of EBUS in diagnosis of bronchogenic cysts [13]. The ultrasonic characteristics of the cyst can be assessed accurately. The EBUS commonly shows the BC as anechoic or hypoechoic well defined structure without color flow signal and differentiates it from tumors or vascular abnormalities. It can also show septations, the presence of hyperechoic thickening within the cyst walls in case of complications [13,14]. It can also examine the echogenicity of the fluid similar to pleural fluids and help assess if it is a source of infection depending on the fluid characteristics [15].

The issues with using EBUS in sampling the structure or in drainage of suspected infected cysts are multifaceted. In the case of a non-complicated cyst, inserting the needle through the bronchoscopic channel and then into the sterile environment of a cyst will surely increase the risk of getting it infected [16]. In the case of an infected bronchogenic cyst, a needle puncture might cause leakage of the infected fluid into the mediastinum, or cause rupture of the cyst with grave consequences [7,11,17]. In both instances, the needle aspiration would not per se prevent re-accumulation of the fluids and thus the recurrence of BC [18]. In addition, It would not eliminate the theoretical increased risk of malignancy since the wall of the cyst persists even though it may not refill with fluids [5]. In a systematic literature review by Maturu et al., 32 patients identified by 26 case reports had EBUS guided drainage of bronchogenic cysts for therapeutic (most commonly) and diagnostic

Download English Version:

<https://daneshyari.com/en/article/8820212>

Download Persian Version:

<https://daneshyari.com/article/8820212>

[Daneshyari.com](https://daneshyari.com)