



Case report

Organized blood clot masquerading as endobronchial tumor: A review of management and recent advances

Lakshmi Saladi^{a,*}, Dmitry Lvovsky^{a,b}^a Division of Pulmonary and Critical Care Medicine, Department of Medicine, Bronx Care Health System, 1650 Grand Concourse, Bronx, NY 10457, USA^b Icahn School of Medicine at Mount Sinai, USA

ARTICLE INFO

Keywords:

Airway obstruction
Blood clot
Bronchoscopy
Cryoextraction

ABSTRACT

Central airway obstruction, a frequently encountered emergency, is usually associated with blood clots, tumors, foreign bodies or mucus plugs. Airway obstruction due to blood clots can be seen as a complication of lung malignancies, infections, bronchiectasis, arteriovenous malformations or pulmonary infarction. In patients with long standing blood clots, the thrombus gets organized and firmly adherent to the airway. The diagnosis is often misleading as these clots mimic tumors clinically and on imaging. Hemoptysis is the most common presenting symptom though many patients can be asymptomatic. Direct visualization with bronchoscopy is required to establish a diagnosis. Life-threatening respiratory impairment is an indication for emergent clot retrieval. Management of these blood clots, especially when organized, is challenging. Initial attempts at removal should include suctioning, lavage or forceps extraction. When unsuccessful, further management options include balloon catheter dislodgement, use of topical thrombolytics, rigid bronchoscopy and cryoextraction.

1. Introduction

Central airway obstruction is a life-threatening emergency and is associated with various conditions like foreign body aspiration, benign or malignant tumors, mucosal edema, impacted mucus and blood clots. Endobronchial obstruction from an organized blood clot mimicking a tumor is unusual, though not rare [1–5]. In the presence of significant hemoptysis, the diagnosis is obvious. However, hemoptysis is not always a presenting feature in these patients and direct visualization with bronchoscopy is required to exclude other conditions [4–6]. The degree of respiratory impairment varies and, sometimes, patients may be asymptomatic. On the other end of the spectrum are those patients who have significant respiratory compromise requiring mechanical ventilation. Managing such patients is challenging and different bronchoscopic modalities like rigid bronchoscopy, Fogarty catheter dislodgement, forceps, snares and cryoextraction were reported to be successful in the past [1,6,11,12,17–24]. However, these modalities are not always readily available and some require expertise. We present a case of an endobronchial organized blood clot that was successfully removed with flexible fiberoptic bronchoscopy using suction and forceps followed by a detailed discussion on the management of such patients.

2. Case presentation

A 54-year-old woman with chronic pulmonary thromboembolism on warfarin, presented to our emergency department with epistaxis and cough with blood tinged sputum of 4-days duration. She had chronic respiratory failure secondary to COPD requiring home oxygen, deep vein thrombosis, hypertension and history of motor vehicle accident many years ago causing chest wall deformities. She required tracheostomy followed by de-cannulation about 20 years prior to current hospitalization. Her family history was significant for lung cancer in her mother. She was an active smoker with more than 25 pack year smoking history.

Physical examination revealed an obese woman in no apparent distress. Her vital signs were stable and chest auscultation revealed bilateral crackles. Laboratory studies were significant for chronic hypercapnia; her hemoglobin was stable. Chest X-ray is shown in Fig. 1.

Antibiotics were initiated for presumed pneumonia and anticoagulation was held. A CT chest was obtained which showed a 5cm long soft tissue lesion in right bronchus intermedius extending into right middle and lower lobes (Fig. 2A and B).

The patient was admitted to the medical floor for further management. Flexible bronchoscopy was performed that revealed a firm, tan colored lesion obstructing 95% of the proximal right mainstem bronchus (Fig. 3).

* Corresponding author.

E-mail addresses: neeha.viks@gmail.com (L. Saladi), dlvovsky@bronxleb.org (D. Lvovsky).

Abbreviations

Chest X-ray	Chest roentgenogram
COPD	Chronic obstructive pulmonary disease
CT chest	Computed tomography of chest
mL	milliliter
U	units

The lesion was noted to have white laminations and an organized thrombus was suspected. The clot was slowly retrieved using suction and biopsy forceps in a piecemeal fashion. Larger pieces of clot were brought out along with the bronchoscope. The lesion extended into right upper, middle and lower lobe bronchi causing complete occlusion. After removal of the extensive organized clot over a span of few hours, no endobronchial lesions were noted (Fig. 4).

There was mild oozing from right lower lobe anterior basal segment. Pulmonary angiogram was performed followed by embolization of the bronchial artery with clinical improvement. Microscopic pathological examination of the lesion confirmed an organized thrombus. She remained clinically stable without recurrence of hemoptysis.

3. Discussion

Airway obstruction from blood clot is a known complication of trauma to tracheobronchial tree, surgical procedures like tracheostomy, alveolar hemorrhage, pulmonary infections like tuberculosis, bronchiectasis, malignancy, pulmonary infarcts and arteriovenous malformations [1,2]. In 1929, Wilson first reported a case of lung atelectasis secondary to a blood clot following hemoptysis in a patient with

pulmonary tuberculosis. Atelectasis improved with spontaneous expectoration of the blood clot [3].

Hemoptysis is a presenting feature in most cases but the diagnosis can be challenging in up to 30% of the cases where it is absent. This is commonly seen in patients on prolonged mechanical ventilation [4–6]. On physical examination, patients present with clinical evidence of atelectasis with decreased breath sounds on the affected side and sometimes a localized wheeze can be heard. Chest X-ray may be completely normal or may show evidence of lung collapse with mediastinal shift to the affected side. Ipsilateral hyperaeration, sometimes leading to pneumothorax, can be seen if the blood clot acts as a one-way valve [7,8]. Direct visualization with bronchoscopy confirms the diagnosis, a fresh blood clot is dark red in color. However, a long-standing organized blood clot appears white or grey in color mimicking a friable tumor or foreign body.

Management of airway obstruction from blood clots is not standardized and depends on the degree of respiratory impairment, hemodynamic stability, site of obstruction and time since obstruction. In a patient who is hemodynamically stable without major respiratory impairment, observation for spontaneous expectoration of the clot is reasonable [2,3,9]. Attempts to remove such clots may lead to further bleeding or more proximal obstruction which can be catastrophic. Freshly formed clots are usually expectorated within 36–48 hours, but may sometimes progress over days and become organized leading to respiratory failure [10]. Both flexible and rigid bronchoscopy are useful in removal of endobronchial blood clots and initial attempts at removing the clot should involve flexible bronchoscopy with saline lavage and suctioning. If unsuccessful, clot retrieval can be performed in a piecemeal manner using biopsy forceps. Larger pieces of clot can be removed by simultaneously withdrawing the forceps and bronchoscope [6].



Fig. 1. Chest X-ray with bilateral opacities and chronic right chest wall deformities.

Download English Version:

<https://daneshyari.com/en/article/8820258>

Download Persian Version:

<https://daneshyari.com/article/8820258>

[Daneshyari.com](https://daneshyari.com)