

Accepted Manuscript

Diagnostic yield of percutaneous biopsy for sclerotic bone lesions: Influence of mean Hounsfield units

Jennifer Ní Mhuirheartaigh, Colm McMahon, Yu-Ching Lin, Jim Wu



PII: S0899-7071(17)30117-1
DOI: doi: [10.1016/j.clinimag.2017.06.008](https://doi.org/10.1016/j.clinimag.2017.06.008)
Reference: JCT 8267

To appear in:

Received date: 28 January 2017
Revised date: 15 June 2017
Accepted date: 28 June 2017

Please cite this article as: Jennifer Ní Mhuirheartaigh, Colm McMahon, Yu-Ching Lin, Jim Wu , Diagnostic yield of percutaneous biopsy for sclerotic bone lesions: Influence of mean Hounsfield units, (2017), doi: [10.1016/j.clinimag.2017.06.008](https://doi.org/10.1016/j.clinimag.2017.06.008)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Diagnostic yield of percutaneous biopsy for sclerotic bone lesions: influence of mean Hounsfield Units

Authors:

1. Jennifer Ní Mhuircheartaigh, MB BCh BAO; Beth Israel Deaconess Medical Center, Harvard Medical School, Department of Radiology
Boston, MA, USA
2. Colm McMahon, MB BCh; Beth Israel Deaconess Medical Center, Harvard Medical School, Department of Radiology
Boston, MA, USA
3. Yu-Ching Lin, MD; Chang Gung Memorial Hospital, Keelung and Chang Gung University, Department of Medical Imaging and Intervention
5 Fu-Shin Street
Kueishan
Taoyuan, TW 333
4. Jim Wu, MD; Beth Israel Deaconess Medical Center, Harvard Medical School, Department of Radiology
Boston, MA, USA

Corresponding author:

Jennifer Ní Mhuircheartaigh

Tel: 617 667 1298

Cell: 617 435 6773

Fax: 617 754 2525

E-mail: jnimhuircheartaigh@hotmail.com

Abstract:

Purpose: To identify whether there was an association between Hounsfield units of sclerotic bone lesions and diagnostic yield of biopsy.

Method: All core needle biopsies of sclerotic bone lesion were identified from a database. Pathology reports were reviewed to determine whether the biopsy was diagnostic or non-diagnostic.

Download English Version:

<https://daneshyari.com/en/article/8821600>

Download Persian Version:

<https://daneshyari.com/article/8821600>

[Daneshyari.com](https://daneshyari.com)