

Accepted Manuscript

Title: Early Onset Inflammatory Bowel Disease – What the Radiologist Needs to Know

Authors: Barber JL, Shah N, Uhlig H, Watson TA

PII: S0720-048X(18)30231-6
DOI: <https://doi.org/10.1016/j.ejrad.2018.07.001>
Reference: EURR 8236

To appear in: *European Journal of Radiology*

Received date: 16-4-2018
Revised date: 15-6-2018
Accepted date: 2-7-2018



Please cite this article as: Barber J, Shah N, Uhlig H, Watson T, Early Onset Inflammatory Bowel Disease – What the Radiologist Needs to Know, *European Journal of Radiology* (2018), <https://doi.org/10.1016/j.ejrad.2018.07.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Early Onset Inflammatory Bowel Disease – What the Radiologist Needs to Know

Barber JL ^{1,3*}, Shah N², Uhlig H ⁴, Watson TA³,

1 Department of Radiology, St George's University Hospitals NHS Foundation Trust, London

2 Department of Paediatric Gastroenterology, Great Ormond Street Hospital, London

3 Department of Radiology, Great Ormond Street Hospital, London

4 Translational Gastroenterology Unit, John Radcliffe Hospital, University of Oxford, Oxford, UK. Department of Paediatrics, University of Oxford, Oxford, UK.

Address correspondence to:

Dr Tom Watson

Department of Radiology

Great Ormond Street Hospital for Children NHS Foundation Trust

London

Wc1N 3JH

tom.watson@gosh.nhs.uk

Type of manuscript: Invited review

Highlights:

- Early-onset bowel disease is defined as disease presenting before the age of 10 years
- Very-early onset disease presents before the age of 6 years and infantile disease presents before 2 years of age
- Very-early onset disease is more likely to be localized to the colon
- There is a strong genetic component to the disease with specific genetic mutations identified in IL-10, XIAP, NCF2, TTC7a, and IPEX

Download English Version:

<https://daneshyari.com/en/article/8822486>

Download Persian Version:

<https://daneshyari.com/article/8822486>

[Daneshyari.com](https://daneshyari.com)