




Chinese Taipei Society of
Ultrasound in Medicine

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.jmu-online.com



ORIGINAL ARTICLE

Assessment of the Anterior Talofibular Ligament Thickness in Patients with Chronic Stroke: An Ultrasonographic Study

Mustafa Turgut Yildizgoren ^{1*}, Onur Velioglu ¹,
Ozcan Demetgul ², Ayse Dicle Turhanoglu ¹

¹ Department of Physical Medicine and Rehabilitation, and ² Department of Neurology, Mustafa Kemal University Medical School, Hatay, Turkey

Received 30 December 2016; accepted 24 January 2017

Available online ■ ■ ■

KEYWORDS

anterior talofibular
ligament,
chronic stroke,
ultrasonography,
walking disabilities

Abstract *Background:* Patients with equinovarus deformity have an increased risk of fall and ankle ligament injury, because of inappropriate prepositioning of the ankle at the end of the swing phase, and inadequate leg and ankle stability during the stance phase. Accordingly, the aim of this study is to compare anterior talofibular ligament (ATFL) thickness of chronic stroke patients with that of healthy individuals using ultrasonography.

Methods: This was a case-control study conducted in a university hospital between July 2015 and July 2016. We included 38 patients [study group; mean age, 59.0 ± 11.1 years; mean body mass index (BMI), 25.4 ± 4.3 kg/m²] and a control group of age-, sex-, and BMI-matched healthy individuals. Demographic and clinical characteristics of the patients (i.e., age, weight, height, Brunnstrom motor recovery stage, Functional Ambulation Scale, Ashworth Scale, and duration of hemiplegia) were recorded during their visits. Furthermore, ultrasound image of the ATFL was obtained from each ankle. The thickness of the ATFL was measured at the midpoint of the ligament between the attachments on the lateral malleolus and the talus using ultrasonography.

Results: In the study group, the mean thickness of the ATFLs of the affected side (2.75 ± 0.41 mm) was thicker than both the unaffected side (2.42 ± 0.30 mm) and the healthy controls (2.35 ± 0.19 mm; $p = 0.007$, $p < 0.001$, respectively). No differences were seen between the two sides of the control group.

Conclusion: Chronic stroke patients have a thicker ATFL on both the affected and unaffected sides, compared with healthy individuals. This architectural feature of the ATFL may be a

Conflicts of interest: The authors have no conflicts of interest to declare.

* Correspondence to: Dr Mustafa Turgut Yildizgoren, Department of Physical Medicine and Rehabilitation, Mustafa Kemal University Medical School, 31060, Serinyol, Hatay, Turkey.

E-mail address: ftm.mustafaturgut@hotmail.com (M.T. Yildizgoren).

<http://dx.doi.org/10.1016/j.jmu.2017.03.001>

0929-6441/© 2017, Elsevier Taiwan LLC and the Chinese Taipei Society of Ultrasound in Medicine. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Yildizgoren MT, et al., Assessment of the Anterior Talofibular Ligament Thickness in Patients with Chronic Stroke: An Ultrasonographic Study, Journal of Medical Ultrasound (2017), <http://dx.doi.org/10.1016/j.jmu.2017.03.001>

result of equinovarus deformity together with spastic muscles. For this reason, early treatment of deformed ligaments and spastic muscles is needed to prevent equinovarus deformity in patients with stroke.

© 2017, Elsevier Taiwan LLC and the Chinese Taipei Society of Ultrasound in Medicine. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Following a stroke event, patients with hemiparesis may develop walking disabilities. On the affected side, muscular imbalance usually causes ankle and foot deformities. Inadequate ankle dorsiflexion and spasticity of the plantar flexors or invertors following stroke have been described [1]. Ankle equinus deformities are seen in up to 20% of stroke patients, and of these, equinovarus deformity is the most characteristic and the most frequently seen [2,3]. Increased tone of the plantar flexors and invertors together with weakness of the dorsal flexors and evertors may explain the development of equinovarus deformity [4,5]. Although hemiplegia is primarily associated with unilateral motor involvement, evident changes in almost all of the parameters used to assess walking have been seen on both the affected and unaffected sides of the body [6].

A ligament is soft connective tissue that transmits tensile force from bone to bone and undertakes an essential role in musculoskeletal biomechanical function by stabilizing and guiding the motion of diarthrodial joints. It tends to remodel according to the applied motion and stress [7,8]. Ligaments supporting the lateral complex of the ankle include the anterior talofibular ligament (ATFL), calcaneofibular ligament, and posterior talofibular ligament. The ATFL is a flat ligament that attaches from the anterior border of the lateral malleolus to the talus, just anterior to the lateral malleolus articular surface. The ATFL limits plantar flexion and inversion, which are the movements coinciding with the most common mechanism of injury [9]. As a result, the ATFL becomes vulnerable in a plantar-flexed and inverted position and is most susceptible to damage during walking in stroke patients. Equinovarus deformity results in an abnormal alignment of the ankle joint, causing the talus to be directed downward and the forefoot to be deviated medially and rotated into supination [10]. Patients with equinovarus deformity have an increased risk of fall and ankle ligament injury, because of inappropriate prepositioning of the ankle at the end of the swing phase, and inadequate leg and ankle stability during the stance phase [11]. However, to the best of our knowledge, the effects of equinovarus deformity on the ATFL have not been investigated in patients with poststroke hemiplegia. Accordingly, the aim of this study was to compare the ATFL thickness in the affected and unaffected sides of chronic stroke patients with that of healthy individuals using ultrasonography.

Materials and Methods

Study design and patients

The study included 38 chronic ischemic or hemorrhagic stroke patients (26 male and 12 female patients) with walking disability. Chronic stroke was defined as the open-ended period starting 6 months after the initial stroke. A control group was formed of 38 sex-, age-, and body mass index (BMI)-matched healthy individuals without ankle disorder or a history of lower limb surgery. Sex, age, BMI, Brunnstrom motor recovery stage, Functional Ambulation Scale (FAS), and duration of hemiplegia were recorded. The Ashworth Scale was used in hemiplegic tonus evaluation. Inclusion criteria were as follows: (1) a stroke event at least 6 months prior to the study and (2) a walking disability with equinovarus of the ankle. Exclusion criteria were as follows: (1) patients with an ankle injury in the poststroke period, (2) previous botulinum toxin injection to gastrosoleus muscles, (3) previous surgery on the lower limbs, (4) fixed ankle contracture, and (5) any reason for pes planus. All patients were informed about the study procedure and informed consent was obtained. The Local Ethics Committee approved the study protocol.

Ultrasonographic measurements

The ATFL measurement was taken using a linear probe (7-12 MHz, Logiq P5, GE Medical Systems, Waukesha, WI, USA) by a physiatrist (M.T.Y.), with the patient in a supine position with the ankles in neutral or slight plantar flexion. A longitudinal image of the ATFL was scanned with the transducer placed in a slightly oblique direction from the anterolateral aspect of the lateral malleolus to the peak of the talus using a previously described method [5]. The peak of the talus represents the anterior aspect of the lateral talar articular cartilage and the lateral neck of the talus. The two bony landmarks were easily identified because of their hyperechogenicity on sonography. The normal ATFL was depicted as hyperechoic bundles on sonography. The thickness of the ATFL was measured halfway between the two bony landmarks of the ankle (Figure 1).

All sonographic measurements were taken in the plane perpendicular to the long axis of the ligaments for standardization and reproducibility of the measurements. Sonographic measurements for the thickness of the ligaments were taken two times, and the mean values were recorded. The reliability of the thickness measurements

Download English Version:

<https://daneshyari.com/en/article/8823512>

Download Persian Version:

<https://daneshyari.com/article/8823512>

[Daneshyari.com](https://daneshyari.com)