

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Interventional Radiology

Skin ischemia and ulceration as a complication of inferior phrenic artery embolization for hepatocellular carcinoma

Sukjin Koh MD, Erik J. Maki MD, Kenneth J. Kolbeck MD, PhD,
Khashayar Farsad MD, PhD*

Charles T. Dotter, Department of Interventional Radiology, Oregon Health and Science University,
3181 SW Sam Jackson Park Road, Portland, OR 97239, USA

ARTICLE INFO

Article history:

Received 22 February 2018

Accepted 28 February 2018

Available online

Keywords:

Transarterial chemoembolization

TACE

Hepatocellular carcinoma

Embolization

ABSTRACT

Nontarget embolization from transarterial liver-directed therapy for hepatocellular carcinoma is a rare complication. We present a case of flank skin ulceration after embolization of the inferior phrenic artery supplying tumor. This complication of inferior phrenic artery embolization underscores caution when using small particles to embolize extrahepatic supply to hepatocellular carcinoma.

© 2018 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Hepatocellular carcinoma (HCC) is the leading primary malignancy of the liver, most often occurring in the setting of chronic liver disease [1]. Transarterial chemoembolization is a proven treatment for HCC, and the primary therapy in the setting of multifocal disease [2]. Additional transarterial treatment for HCC includes bland embolization, embolization using drug-eluting microspheres, and radioembolization using beta-emitting microspheres [3–5]. Extrahepatic vessels can often supply intrahepatic tumors, particularly those along the capsular surface [6,7]. The right inferior phrenic artery, for example, can supply tumor at the dome or lateral margins of the liver [8,9]. Extrahepatic vasculature can be targeted for transarterial treatment of liver tumors. With any transarterial therapy, nontarget

embolization is a potential complication either by direct or collateral flow. In rare cases, nontarget embolization has resulted in gastrointestinal ulceration, pulmonary infarction, or skin ischemia [10–13].

Before treatment with yttrium-90 (Y90) microspheres, a planning angiogram is done to map out the arterial anatomy of the liver and to calculate the hepatopulmonary shunt fraction using microaggregated albumin Tc99m. On occasion, if extrahepatic vessels are seen to supply the liver tumors, embolization of these vessels may be performed to facilitate Y90 delivery via intrahepatic arteries [13,14]. Additionally, embolization of extrahepatic arterial supply to the tumors can help ensure a complete treatment by treating all pertinent arterial feeders. We present a case of flank skin ischemia and ulceration after transarterial embolization of the right inferior phrenic artery using small particles to address extrahepatic arterial

Competing Interests: Authors have no conflicts of interest to disclose.

* Corresponding author.

E-mail address: farsad@ohsu.edu (K. Farsad).

<https://doi.org/10.1016/j.radcr.2018.02.032>

1930-0433/© 2018 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

supply during a planning angiogram for radioembolization. The danger of small particle embolization of extrahepatic feeders to HCC leading to nontarget ischemic effects is underscored.

Case presentation

Institutional review board approval is not required for single patient retrospective review at our institution. A 66-year-old male with hepatitis C virus-related cirrhosis and portal hypertension was referred for treatment of an infiltrative 10-cm right hepatic lobe HCC (Fig. 1A). The patient was selected for Y90 radioembolization based on consensus recommendations by our multidisciplinary liver tumor board. The patient's liver function was preserved (Child-Pugh-Turcotte score A5), and he had good functional status with no history of ascites, hepatic encephalopathy, or gastrointestinal bleeding.

Planning angiography demonstrated dominant tumor arterial supply from the posterior division of the right hepatic artery (Fig. 1B). In addition, angiography of the right inferior phrenic artery showed approximately 10% supply to the overall tumor burden (Fig. 1C). Particle embolization of the right inferior phrenic supply was performed at the time of planning angiography with plans for subsequent Y90 treatment to the dominant hepatic arterial supply. One-hundred micron microspheres were used to embolize the right inferior phrenic artery vascular supply to the right lobe HCC. Postembolization

angiography showed decreased tumor enhancement and preserved antegrade flow in the right inferior phrenic artery (Fig. 1D).

Approximately 8 hours after treatment, the patient reported right flank pain and had slight mottling to the skin in a dermatomal pattern (Fig. 2A). The skin changes worsened over the ensuing several hours, prompting a punch biopsy of the affected area. The biopsy specimen revealed microspheres within the vasculature, suggesting ischemic dermatopathy as the cause of the observed skin changes (Fig. 3). The patient was managed conservatively with emollient application to the affected area for moist healing, and was given oral narcotic pain medication for associated discomfort. The skin was monitored for the next 3 weeks, during which time 2 focal areas of skin ulceration measuring approximately 1 cm each were observed (Fig. 2B). Conservative management continued, and his skin healed over the next 3 months without progressive ulceration or infectious complication, leaving only focal residual pigmentation (Fig. 2C).

Discussion

Nontarget embolization is a rare complication of transarterial therapies for hepatic neoplasms [10,13]. Particularly when treating intrahepatic tumors via extrahepatic collateral circulation, caution must be diligently applied to avoid embolization of anatomic perfusion pathways. Structures at risk for such nontarget




Fig. 1 – (A) Arterial phase MRI showing an infiltrative arterial enhancing mass in the posterior right lobe of the liver. (B) Angiogram from the right posterior division hepatic artery showing hypertrophy of the artery and dominant supply to the right lobe HCC. (C) Angiogram from the right inferior phrenic artery showing supply to the superior and medial aspects of the tumor (arrows). (D) Postembolization angiogram showing decreased tumor blush supplied by the right inferior phrenic artery. HCC, hepatocellular carcinoma; MRI, magnetic resonance imaging.

Download English Version:

<https://daneshyari.com/en/article/8825027>

Download Persian Version:

<https://daneshyari.com/article/8825027>

[Daneshyari.com](https://daneshyari.com)