

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Interventional Radiology

Variceal bleeding in the small intestine successfully treated with balloon-occluded retrograde transvenous obliteration using N-butyl-2-cyanoacrylate: A case report

Koichi Masuda MD^{a,*}, Shinsuke Takenaga MD^a, Hirokazu Ashida MD^b,
Hiroya Ojiri PhD, MD^b

^a Department of Radiology, The Jikei University Katsushika Medical Center, 6-41-2, Aoto, Katsushika-ku, Tokyo, 125-8506, Japan

^b Department of Radiology, The Jikei Medical University Hospital, Minato-ku, Tokyo, Japan

ARTICLE INFO

Article history:

Received 30 October 2017

Accepted 15 January 2018

Available online 6 February 2018

Keywords:

Small intestinal varices

Balloon-occluded retrograde

transvenous obliteration

N-butyl-2-cyanoacrylate

ABSTRACT

Rupture of small intestinal varices associated with portal hypertension can be a serious condition that is difficult to diagnose early and to manage. Moreover, optimal guidelines for the treatment of small intestinal varices have not yet been established. We herein report a case of a 73-year-old man with small intestinal varices. The man presented with bleeding from a stoma in the small intestine, which subsequently led to hemorrhagic shock. We successfully treated the patient with balloon-occluded retrograde transvenous obliteration via the right inferior epigastric vein using N-butyl-2-cyanoacrylate.

© 2018 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Ectopic varices, caused by portal hypertension, are defined as portosystemic collaterals located outside the gastroesophageal region [1]. Ectopic varices can cause gastrointestinal bleeding and can be life threatening. However, ectopic varices are difficult to diagnose, manage, and treat because there are no optimal guidelines for the treatment of small intestinal varices currently.

Balloon-occluded retrograde transvenous obliteration (B-RTO) is an angiographic technique that has been accepted as a treatment option for small intestinal varices [2]. B-RTO using N-butyl-2-cyanoacrylate (NBCA; Histoacryl; B. Braun, Melsungen, Germany), a liquid embolic material that works rapidly and permanently [3], has been reported to be useful for the treatment of bleeding duodenal varices [4].

We herein report a case regarding the successful treatment of ruptured small intestinal varices successfully treated by B-RTO via the right inferior epigastric vein using NBCA.

Competing Interests: All authors declare that they have no conflicts of interest.

* Corresponding author.

E-mail address: masu0205@icloud.com (K. Masuda).

<https://doi.org/10.1016/j.radcr.2018.01.023>

1930-0433/© 2018 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – (A) Contrast-enhanced computed tomography scan reveals small intestinal varices and extravasation (arrow). (B) Three-dimensional computed tomography scan waving toward the right anterior oblique position reveals small intestinal varices (arrowhead), as well as the branch vessel of the superior mesenteric vein (arrow) and the right inferior epigastric vein (open arrow).

Case report

Case

A 73-year-old man presented to our hospital with bleeding from an ileostomy. The man had suffered from liver cirrhosis caused by nonalcoholic steatohepatitis and had undergone hemicolectomy and ileostomy because of ascending colon cancer 18 months previously. The patient's initial laboratory data were as follows: red blood cell count, $2.60 \times 10^6/L$; hemoglobin, 8.8 mg/dL (anemia); platelet count, $107 \times 10^3 \mu L$; total bilirubin, 0.8 mg/dL; albumin, 3.0 g/dL; fibrinogen, 199 mg/dL; and international normalized ratio of prothrombin time, 1.45. Six months previously, an upper gastrointestinal endoscopy was performed, and only scarring from esophageal varices was detected.

Emergency endoscopy from the ileostomy could not reveal the bleeding point. Moreover, a gradual drop in blood pressure was observed (systolic blood pressure, 60 mm Hg). To determine the source of bleeding, urgent contrast-enhanced computed tomography (CT) was performed. The CT showed small intestinal varices with extravasation and a small amount of ascites (Fig. 1).

Therefore, emergency angiography was performed. The right common femoral vein was punctured under local anesthesia, and a 6-Fr sheath introducer was inserted. Subsequently, a 4-Fr cobra-shaped catheter was inserted into the right inferior epigastric vein. Using coaxial technique, a 2.7-Fr microballoon catheter (Attendant Δ ; Terumo Clinical Supply, Gifu, Japan) was advanced into the proximal portion of the small intestinal varices. Balloon-occluded retrograde transvenous venography via the right inferior epigastric vein revealed small intestinal varices without extravasation (Fig. 2).

Complete variceal embolization was achieved using a 0.5 mL NBCA : lipiodol mixture (1:1) (Fig. 3). The duration from the injection of the NBCA : lipiodol mixture until balloon deflation was approximately 20 seconds. After embolization, the patient

had no additional bleeding and was discharged 20 days after B-RTO.

Discussion

Varices of the esophagus and the stomach are common portosystemic collaterals in patients with portal hypertension; however, ectopic varices are uncommon. Approximately 17% of ectopic varices are found in the jejunum and the ileum, 17% are found in the duodenum, 14% in the colon, 8% in the




Fig. 2 – The right inferior epigastric vein was the main draining vein. Balloon-occluded retrograde transvenous venography via the right inferior epigastric vein shows small intestinal varices (arrow) with no extravasation.

Download English Version:

<https://daneshyari.com/en/article/8825104>

Download Persian Version:

<https://daneshyari.com/article/8825104>

[Daneshyari.com](https://daneshyari.com)