

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Case Report

Bow hunter's syndrome in a patient with vertebral artery atresia, an arcuate foramen, and unilateral deafness: a case report

Charles T. Simpkin MS^{a,*}, Kelly E. Davis^a, Brady S. Davis DO^b, Andrew M. Vosko PhD^a, Michael E. Jorgensen PhD^a

^a Department of Structural Medicine, Rocky Vista University, College of Osteopathic Medicine, 8401 S Chambers Rd, Parker, CO 80134, USA

^b Department of Radiology, David Grant USAF Medical Center, Travis Air Force Base, CA, USA

ARTICLE INFO

Article history:

Received 24 March 2017

Accepted 4 April 2017

Available online xxx

Keywords:

Bow hunter's syndrome

Vertebral artery atresia

Arcuate foramen

Sensorineural deafness

ABSTRACT

Bow hunter's syndrome (BHS) is a rare cause of vertebrobasilar insufficiency that occurs when the vertebral artery (VA) is occluded on rotation of the head and neck. This dynamic occlusion of the VA can occur anywhere along its course after it arises from the subclavian artery. Although most cases are associated with compression by osteophytes, cervical spondylosis, or lateral disc herniation, BHS has a highly variable clinical course that depends on the patient's specific anatomy. Therefore, it may be important for clinicians to be aware of anatomical variants that predispose individuals to BHS. Here, we report on a patient with BHS who was found to have two uncommon anatomical anomalies: an atretic right VA and a left-sided arcuate foramen.

© 2017 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Structures within the posterior cranial fossa, including the brainstem and cerebellum, are predominantly vascularized by the vertebral arteries (VAs). Ischemic events involving this posterior circulation account for 20% of all strokes and can result in devastating paralysis and even death [1]. Transient or

incomplete ischemia within this vascular territory can cause vertebrobasilar insufficiency (VBI), which is characterized by symptoms such as vertigo, nystagmus, and syncope [2]. Rarely, VBI can be transiently induced by rotation of the head and neck [3]. This condition is known as Bow hunter's syndrome (BHS) or rotational VA occlusion. BHS was first formally described in 1978 by B. F. Sorensen in a patient who suffered

Acknowledgements: We would like to extend our thanks to the patient for allowing us to share her case. Also to the Pre-Doctoral Anatomy Fellowship at Rocky Vista University College of Osteopathic Medicine which provided the opportunity and resources to complete this study.

Informed Consent: Written informed consent for experimentation with human subjects was obtained from the patient.

Approved by Rocky Vista University IRB: HRIB 2016-00003.

Competing Interests: The authors have declared that no competing interests exist.

* Corresponding author.

E-mail address: charles.simpkin@rvu.edu (C.T. Simpkin).

<http://dx.doi.org/10.1016/j.radcr.2017.04.001>

1930-0433/© 2017 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Wallenberg syndrome after archery practice [4]. Although this initial case report described a patient who suffered permanent deficits after a rotation-induced occlusive event, the term BHS is now broadly used to describe a reversible condition.

The mechanism underlying typical BHS can be conceptualized as a “two-hit” model. The “first-hit” may be congenital or acquired, most often one VA is severely hypoplastic or stenotic. The “second-hit” occurs when the contralateral (and now dominant) VA is occluded by a bony-overgrowth, fibrous band, or herniated disc on rotation of the head [5,6]. Atlantoaxial compression of the dominant VA by a cervical spine osteophyte has been shown to be the most common cause of BHS [7]. However, this model is limited in describing all cases. The VA can be compressed anywhere along its course into the posterior cranial fossa. There are even case-reports documenting BHS after the occlusion of the non-dominant VA [8]. As a result, the clinical presentation and radiographic findings in patients with BHS are highly variable [3]. A more thorough understanding of both normal and abnormal VA anatomy may be helpful for clinicians while evaluating patients who present with symptoms of VBI. Here, we present a case of BHS in a patient with alterations in both the course and structure of the VAs.

Case report

A 59-year-old Caucasian female presented to her pain management physician for a preprocedural evaluation. She was scheduled to undergo a cervical facet rhizotomy, but on questioning the patient reported dizziness when she rotated her head to the right. These episodes lasted only a few seconds and were associated with lightheadedness and nausea. All symptoms resolved completely on returning her head to the neutral position. The patient reported that these symptoms had started within the past year, but they did not significantly affect her daily life. The dizziness could not be reproduced with any other head or neck movements. There was no recent history of illness, vomiting, tinnitus, changes in vision, postural instability, or syncopal episodes. Her medical history was significant for chronic bilateral neck and shoulder pain of 13 years and right-sided congenital sensorineural deafness. Her hearing loss was diagnosed at around age 3 years without apparent cause. She was not aware of any significant childhood illnesses, head trauma, exposure to ototoxic drugs, or a family history of hearing loss.

As a result of the patient's report of rotational dizziness, a magnetic resonance angiogram was ordered to rule out any major vascular occlusion or dissection. Axial time-of-flight noncontrast magnetic resonance angiogram demonstrated normal right and left internal carotid arteries. The left VA was of normal course and caliber. There was reduced flow through the right VA, which was suspicious for a VA dissection (Fig. 1). The patient was promptly sent to the emergency department for confirmatory computed tomography angiography. These images demonstrated a dominant left VA, which was the sole supply to the basilar artery. The V1-V4 segments of the right VA were markedly hypoplastic, but patent into the posterior cranial fossa. The V4 segment terminated as the right posterior inferior cerebellar artery and did not contribute toward the




Fig. 1 – Posterior view of 3D time-of-flight magnetic resonance angiogram (MRA) rendering. The right vertebral artery is decreased in caliber (arrow). The left vertebral artery (arrowhead) and basilar artery (double arrows) are both of normal course and caliber.

basilar artery (Fig. 2). On the left side of C1, a bony bridge can be seen extending from the lateral mass, over the groove for the left VA, and nearly connecting to the posterior arch (Fig. 3). There was no further evidence of dissection or infarction, so the patient was discharged from the emergency department and received the facet rhizotomy several days later.

Discussion

This case highlights several interesting variations in the course and structure of the VA. Upward of 69% of healthy adults are thought to have some degree of size discrepancy between the right and left VAs, with the left being larger than the right in most cases [9–11]. Vertebral artery hypoplasia (VAH) is a condition that occurs when one VA is significantly smaller than the other, but still connects to the basilar artery system [12]. The diameter cut-off used to define VAH varies depending on the study, but generally range from <2 to <3 mm [13]. VAH has been reported in anywhere from 2.34%–28.9% of healthy adults, with the right VA more frequently affected than the left [14–16]. Although most cases of VAH are asymptomatic, numerous studies have identified it as an independent risk factor for regional hypoperfusion and posterior circulation infarction [13,17,18].

The computed tomography angiography findings in our patient revealed a hypoplastic VA that terminated as the posterior inferior cerebellar artery instead of the basilar artery, which meets criteria for the diagnosis of vertebral artery

Download English Version:

<https://daneshyari.com/en/article/8825365>

Download Persian Version:

<https://daneshyari.com/article/8825365>

[Daneshyari.com](https://daneshyari.com)