

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Case Report

Carbon dioxide angiography and arterial embolization could successfully control postpartum uterine hemorrhage for the patient with hypersensitivity to iodine compound

Sayaka Kawashita MD^{a,b}, Masanori Kaneuchi MD, PhD^{a,*}, Daisuke Nakayama MD, PhD^b, Naoki Fuchi MD^a, Yuko Murakami MD^a, Kiyonori Miura MD, PhD^a, Hideaki Masuzaki MD, PhD^a

^a Department of Obstetrics and Gynecology, Nagasaki University School of Medicine, 7-1, Sakamoto 1-choume, Nagasaki-city 8528501, Japan

^b Department of Obstetrics and Gynecology, Sasebo City General Hospital, Sasebo-city 8578511, Japan

ARTICLE INFO

Article history:

Received 9 December 2016

Received in revised form

16 January 2017

Accepted 22 January 2017

Available online xxx

Keywords:

Postpartum hemorrhage

Carbon dioxide

Iodine allergy

ABSTRACT

Transcatheter arterial embolization has become a common management option for intractable postpartum uterine hemorrhage. Iodinated contrast agents are commonly used for embolotherapy but cannot be used in patients with renal dysfunction or iodine allergy. A 37-year-old woman spontaneously delivered a healthy girl but developed severe postpartum uterine hemorrhage, probably due to placenta accrete. Contrast-enhanced computed tomography revealed extravasation of the iodinated contrast agent into the uterine cavity. The patient subsequently developed allergic reaction to this agent. After recovery from the allergic reaction with appropriate treatment, we performed carbon dioxide angiography and found that the extravasation originated from the right uterine artery. Embolization of the right internal iliac artery was performed, and hemostasis was verified. Based on this experience, we suggest carbon dioxide angiography and arterial embolization can be used for treating intractable postpartum hemorrhage in patients with iodinated contrast media allergy or renal dysfunction.

© 2017 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Postpartum uterine hemorrhage is an obstetric emergency and remains a major cause of maternal deaths worldwide. Primary management options for this condition include bimanual uterine massage, uterotonic agent administration, and balloon tamponade. Another reliable and safe

alternative is transcatheter arterial embolization, which has also proven effective for intractable postpartum uterine hemorrhage [1], alongside other surgical approaches such as compression sutures, ligation of responsible artery, and hysterectomy. Indeed, arterial embolization is included in management guidelines for critical bleeding in obstetrics [2].

Competing Interests: The authors have declared that no competing interests exist.

* Corresponding author.

E-mail address: mkaneuc@hotmail.com (M. Kaneuchi).

<http://dx.doi.org/10.1016/j.radcr.2017.01.017>

1930-0433/© 2017 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Transcatheter arterial embolization is highly effective at achieving hemostasis and recently has helped to reduce the number of patients requiring hysterectomy. An iodinated contrast agent is typically used for angiography and arterial embolization, but caution must be taken when using such agents due to the possibility of an allergic reaction and renal toxicity. Mild allergic symptoms include urticaria, nausea, and vomiting, while serious anaphylactic reaction involving constriction of the airways, low blood pressure, and arrhythmia may be life threatening. In Japan, those serious reactions to iodinated contrast media was observed in 0.04%–0.22% [3]. In patients with a history of iodine allergy or renal dysfunction, carbon dioxide (CO₂) is occasionally used as an alternative contrast media [4].

We present here a case of a woman with severe postpartum uterine hemorrhage who developed an allergic reaction to the iodinated contrast agent during contrast-enhanced computed tomography (CT), which we were able to control successfully by using CO₂ as an alternative contrast medium for angiography and arterial embolization.

Case

A 37-year-old nulliparous woman vaginally delivered a healthy girl weighing 3460 g at 39 weeks and 5 days of gestation in a local hospital. Without spontaneous expulsion, the placenta was manually removed and a small defect was noted. Hemorrhage from the uterine cavity persisted, with more than 700 mL of blood loss occurring within a 30-minute period after removal of the placenta. The patient showed signs of hemorrhagic shock and was urgently transported to our hospital.

On arrival, her blood pressure and heart rate were 74/40 mm Hg and 110 beats/min, respectively. Laboratory data indicated anemia (erythrocyte count $298 \times 10^4/\mu\text{L}$, hemoglobin 9.7 g/dL, hematocrit 31%) and an elevated fibrin degradation product level (22.0 $\mu\text{g/mL}$). Her obstetric disseminated intravascular coagulation score [5] was 8. Ultrasound imaging showed that part of the placenta had remained in the uterine cavity. Upon manually examining the uterine cavity, we removed a 5-cm piece of the placenta. Administration of oxytocin decreased the hemorrhagic flow with satisfactory uterine contraction. Simultaneously, we initiated transfusion of packed red blood cells and fresh frozen plasma and administered gabexate mesilate (1500 mg/d), urinastatin (100,000 IU/d), and antithrombin III (3000 IU/d) to treat disseminated intravascular coagulation. During the initial 12 hours after her arrival at our hospital, 20 units of red blood cell and 41 units of fresh frozen plasma were needed to maintain hemodynamics. Through these intensive treatments, her physical condition improved temporarily.

However, 8 hours after the full placental expulsion was completed at our hospital, uterine hemorrhage increased again and blood loss >1000 mL was observed within a 1-hour period. Contrast-enhanced CT was performed to locate the source of the hemorrhage. After administration of the iodinated contrast agent, iopamidol which was one of the low-osmolar iodinated contrast media (LOCM), the patient experienced flushing of the skin, coughing, and nausea. Wheezing could also be heard over



Fig. 1 – Contrast-enhanced computed tomography showed extravasation of the iodinated contrast agent into the uterine cavity (arrow).

both lungs. These symptoms disappeared after intramuscular injection of adrenaline (0.3 mg) and intravenous injection of methylprednisolone (125 mg).

We hypothesized that the patient had experienced an allergic reaction to the iodinated contrast agent. The limited images obtained by contrast-enhanced CT showed extravasation of the agent into the uterine cavity (Fig. 1). Therefore, we considered that arterial embolization or surgery was required to achieve hemostasis. Since the patient was a primipara and had a strong desire to preserve her uterus, we decided to perform angiography using CO₂ instead of an iodinated contrast agent and arterial embolization.

A 4-Fr sheath was placed in the left femoral artery, and a catheter tip was then placed in the right internal iliac artery. Contrast imaging with CO₂ confirmed contrast leakage through the right uterine artery (Fig. 2A). A microcatheter was inserted in the internal iliac artery, and embolization was performed using pieces of gelatin sponge. Subsequently, hemostasis was verified (Fig. 2B). A total of 80 mL of CO₂ was used without any complications.

Ten days after this treatment, she was discharged well. Pituitary hormone levels were within normal ranges at 1 month postpartum. Magnetic resonance imaging at 2 months postpartum showed no unusual findings in the uterus, and normal menses resumed at 6 months postpartum.

Discussion

We successfully used CO₂ in the present patient who was not known to be allergic to iodinated contrast media until such an agent was used in the management of severe postpartum uterine hemorrhage. To the best of our knowledge, this is the first case report describing the use of CO₂ angiography and arterial embolization for postpartum uterine hemorrhage.

Download English Version:

<https://daneshyari.com/en/article/8825401>

Download Persian Version:

<https://daneshyari.com/article/8825401>

[Daneshyari.com](https://daneshyari.com)