

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

Case Report

High-grade transitional cell carcinoma masquerading as a xanthogranulomatous pyelonephritis and perinephric abscess

Flavio V. Ordones MD^{a,b,*}, Krishanu Das FRCS^a, Simon Prowse MBBS, FRCR, FRANZCR^c, Penelope Cohen FRCPA^d, Nicholas R. Brook BSc, MSc, BM, FRCS(Urol), MD, FRACS(Urol)^a

^a Department of Urology, Royal Adelaide Hospital, North Terrace, Adelaide, SA 5000, Australia

^b Division of Urology, University of São Paulo Medical School, HCFMUSP – Scholarship João Alves de Q. Filho, Sao Paulo, SP, Brazil

^c Department of Radiology, Royal Adelaide Hospital, Adelaide, Australia

^d Department of Pathology, Royal Adelaide Hospital, Adelaide, Australia

ARTICLE INFO

Article history:

Received 3 October 2016

Received in revised form

6 November 2016

Accepted 23 November 2016

Available online xxx

Keywords:

Transitional cell carcinoma

Xanthogranulomatous

Pyelonephritis

ABSTRACT

Xanthogranulomatous pyelonephritis (XGPN) is an atypical long-term pyelonephritis with destruction of renal parenchyma and a long-term inflammatory infiltrate of macrophages. Reported presentations of transitional cell carcinoma (TCC) are different. A 73-year-old woman presented with loin pain, prostration, and fever. Computed tomography scan revealed poor cortical enhancement of the kidney, but some of the images bore resemblance to the characteristic “bear’s paw” sign, consistent with XGPN with a 7-cm perinephric collection. She was provisionally diagnosed as severe acute pyelonephritis, possibly XGPN, with abscess. In view of the poor clinical condition, decision was made to perform nephrectomy. Histology revealed a G3pT4 high grade TCC with perineural and vascular invasion and reactive xanthogranulomatous inflammatory response. There are few reports of concomitant XGPN and TCC affecting the kidney. However, there has not been any mention of XGPN and TCC presenting as acute pyelonephritis and perinephric abscess so far.

Crown Copyright © 2016 Published by Elsevier Inc. on behalf of under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Xanthogranulomatous pyelonephritis (XGPN) is an atypical form of long-term pyelonephritis, characterized by the destruction of renal parenchyma and replacement with a

long-term inflammatory infiltrate of macrophages, known as xanthoma cells. Commonly reported presentations of transitional cell carcinoma (TCC) arising from renal pelvis are macroscopic or microscopic hematuria, flank pain, renal or clot colic, mass, constitutional symptoms, or occasionally

Authors' contributions: FVO, Urology Fellow, Royal Adelaide Hospital, manuscript and references; KD, Urology Fellow, Royal Adelaide Hospital, manuscript and references; SP, Radiology Department, Royal Adelaide Hospital, radiology images; PC, Pathology Department, Royal Adelaide Hospital, pathology report, micro- and macropathology images; NRB, Consultant Urology, Royal Adelaide Hospital, senior author and manuscript editor.

Acknowledgments: Thanks to Dr Penny Cohen for providing the macro- and micropathological images.

Competing Interests: The authors have declared that no competing interests exist.

* Corresponding author.

E-mail address: fvordones@gmail.com (F.V. Ordones).

<http://dx.doi.org/10.1016/j.radcr.2016.11.018>

1930-0433/Crown Copyright © 2016 Published by Elsevier Inc. on behalf of under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – (A) Coronal computed tomography (CT) image at presentation showing perinephric and perisplenic collection with “bear’s paw” sign of the left kidney. (B) Axial CT image after percutaneous drainage with nonresolution of perinephric collection. (C) DMSA renal scan shows poor tracer uptake in left kidney consistent with poor function. DMSA, dimercaptosuccinic acid.

incidental diagnosis from radiological investigations for other pathologies. We report an unusual case of high-grade TCC presenting with a clinical and pathological presentation of XGPN and perinephric abscess.

Case report

A 73-year-old woman presented as an emergency with left loin pain, prostration, and fever. The history revealed of recurrent *Escherichia coli* urinary tract infection (UTI) over the preceding few months, that was treated with multiple courses of antibiotics by her primary care physician. Physical examination revealed tachycardia (110/min), flushed facies, and a distended abdomen with left flank tenderness, but no guarding or peritonitis. Serum parameters were suggestive of acute infection with a neutrophilia ($23.3 \times 10^9/L$), mild anemia (hemoglobin 98 g/dL), and elevated C-reactive protein levels (400 mg/dL). Computed tomography scan with intravenous contrast revealed poor cortical enhancement of the left kidney, but some of the images bore resemblance to the characteristic “bear’s paw” sign, consistent with XGPN. A 7 cm perinephric and perisplenic collection with gas locules posterolateral to the left kidney, and multiple small para aortic nodes were also noted (Fig. 1A.). Chest x-ray was suggestive of left lower lobe consolidation and pleural effusion.

Case hypothesis

She was given a provisional diagnosis of severe acute pyelonephritis, possibly XGPN, with perinephric abscess. Parenteral

antibiotics (ceftriaxone and metronidazole) were initiated with supportive care. A percutaneous drain was inserted under computed tomography guidance into the perinephric collection (Fig. 1B). Purulent fluid was drained, from which *E. coli* infection was cultured. Despite optimal supportive measures, only marginal clinical improvement was noted over the next few days. She continued to experience high grade fever, flank pain, leukocytosis, and serum acute phase reactants continued to rise. Despite repeated saline flushes, output from the drain was low and repeat imaging revealed unresolved pyelonephritis and perinephric and peri splenic collection. In view of the continuing poor clinical condition, a decision was made to perform nephrectomy and thorough lavage of perinephric and perisplenic area. This was further supported by the poor function (11%) of the left kidney on dimercaptosuccinic acid renogram (Fig. 1C.). Nephrectomy and lavage was undertaken through a retroperitoneal approach. The procedure and postoperative recovery were uneventful.

Histology revealed a G3pT4 high-grade TCC traversing the cortex and medulla with capsular penetration, perineural and vascular invasion, and a xanthogranulomatous inflammatory stromal response (Figs. 2A and B). Resection margins were free of tumor.

Discussion

Acute presentation with fever, loin pain, hemodynamic instability, and elevated acute phase reactants on the background of recurrent UTI is not typical of upper tract TCC. The presentation led us to consider severe acute pyelonephritis (with perinephric abscess) as a provisional diagnosis. Imaging also pointed to aggressive infection with pus. The usual

Download English Version:

<https://daneshyari.com/en/article/8825408>

Download Persian Version:

<https://daneshyari.com/article/8825408>

[Daneshyari.com](https://daneshyari.com)