

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: <http://Elsevier.com/locate/radcr>

## Case Report

# Beware of positive pressure: coronary artery air embolism following percutaneous lung biopsy

Dmitriy N. Kazimirko MD<sup>a,\*</sup>, William B. Beam MD<sup>b</sup>, Khalid Saleh MD<sup>a</sup>, Akash M. Patel MD<sup>a</sup>

<sup>a</sup> Department of Radiology, University of Mississippi Medical Center, 2500 N State St, Jackson, MS 39216, USA

<sup>b</sup> Department of Anesthesiology, Mayo Clinic, 200 First St SW, Rochester, MN, 55905, USA

### ARTICLE INFO

#### Article history:

Received 8 May 2016

Received in revised form

18 July 2016

Accepted 12 August 2016

Available online xxx

#### Keywords:

Systemic air embolism

Coronary artery air embolism

Percutaneous lung biopsy

Transient heart block

### ABSTRACT

Patients undergoing percutaneous lung biopsy are at risk of developing a systemic air embolism. Air embolism may manifest as a catastrophic iatrogenic event with ischemic insult to the end organs, with sites of least resistance such as coronary and cerebral circulation the most susceptible. We review the available literature and present a case of iatrogenic air embolism during computed tomography guided percutaneous lung biopsy under general anesthesia. Management, outcome, and periprocedural factors that may have contributed to the complication are discussed.

© 2016 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Computed tomography (CT) guided percutaneous lung lesion biopsy is an indispensable diagnostic tool. Complications related to this procedure include pneumothorax, pulmonary hemorrhage, tumor seeding, and systemic air embolism (SAE), with latter reported incidence of 0.06% [1]. SAE may manifest as a catastrophic iatrogenic event with ischemic insult to the end-organ sites of least resistance, such as coronary or cerebral circulation. We present a case of a patient who suffered an iatrogenic coronary air embolism during CT-guided percutaneous lung biopsy manifesting as a heart block.

## Case report

A 65-year-old man referred for biopsy of a right middle lobe nodule. Because of the proximity of the nodule to the diaphragm and major fissure, the case was performed with patient positioned right side up 30° and anesthesia with endotracheal positive airway pressure to facilitate an optimal access window by hyperexpanding the lung and controlling respiration (Fig. 1).

A 20-gauge coaxial system was directed into the lesion followed by 24-gauge coaxial needle aspirates. Attempts to redirect the access needle into the lesion resulted in moderate iatrogenic pneumothorax (Fig. 2A) necessitating placement of an 8-French pigtail catheter connected to wall suction into the pleural space

Competing Interests: The authors have declared that no competing interests exist.

\* Corresponding author.


E-mail address: [dkazimirko@umc.edu](mailto:dkazimirko@umc.edu) (D.N. Kazimirko).

<http://dx.doi.org/10.1016/j.radcr.2016.08.007>

1930-0433/© 2016 the Authors. Published by Elsevier Inc. under copyright license from the University of Washington. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



**Fig. 1 – Coronal contrast enhanced CT chest. Right middle lobe lung nodule, abutting the diaphragm.**



**Fig. 3 – Axial nonenhanced CT chest. Air-blood level in the superior pulmonary vein and ascending aorta post biopsy.**


with near complete resolution of pneumothorax (Fig. 2B). Procedure was completed after 5 fine-needle aspirates.

At the completion of sample collection, telemetry revealed intermittent third degree heart block with junctional rhythm in the 30 bpm range (telemetry strip unavailable). Transcutaneous pacing was instituted with capture at a rate of 60 bpm. Delivered oxygen was increased to 100%, and the table was placed in Trendelenburg. With these interventions, the patient remained hemodynamically stable without the need for additional supportive therapies.

Postprocedural CT scan following needle removal revealed iatrogenic air embolism in the segmental pulmonary vein draining the biopsy site (Fig. 2C) with nondependent air collecting in the superior right pulmonary vein (2D) and aorta (Figs. 2D and 3). Additionally, air was noted in the entirety of the right coronary artery (RCA), conus branch, proximal, mid,

and distal RCA including 2 acute marginal, posterior descending, and posterolateral branches (Figs 4A-E). Heart block was likely a manifestation of air embolism to the atrioventricular nodal branch. Ten-minute and 20-minute delayed CT scans of the chest demonstrated gradual decrease with complete resolution of air contained within the aorta and coronary arteries 40 minutes later. CT imaging of the head did not reveal gas in the intracranial vasculature.

With resolution of intracoronary air, the rhythm returned back to normal sinus, and transcutaneous pacing was discontinued. General anesthesia was discontinued, and the patient was extubated. On examination, patients only complaint was reproducible pain at the chest tube site. He denied other chest pain sensation, headache, abdominal pain, weakness, and confusion. Follow-up 12-lead electrocardiogram showed borderline (mild prolongation of PR interval to 220 ms) first



**Fig. 2 – (A-D). CT-guided nodule biopsy. Iatrogenic pneumothorax (A), with resolution after ventral pigtail placement, (B). Air filling a segmental pulmonary vein next to a bronchus (red circle, C), which is draining into a nondependent superior pulmonary vein and embolized to the ascending aorta (red arrows, D).**

Download English Version:

<https://daneshyari.com/en/article/8825545>

Download Persian Version:

<https://daneshyari.com/article/8825545>

[Daneshyari.com](https://daneshyari.com)