

Author's Accepted Manuscript

Risk Factors for Readmission Following Shock Wave Lithotripsy

Benjamin Nelson, MD, Anees Fazili, MD, Erdal Erturk, MD



PII: S2352-0779(16)30175-3
DOI: [10.1016/j.urpr.2016.04.002](https://doi.org/10.1016/j.urpr.2016.04.002)
Reference: URPR 199

To appear in: *Urology Practice*
Accepted Date: 30 April 2016

Please cite this article as: Nelson B, Fazili A, Erturk E, Risk Factors for Readmission Following Shock Wave Lithotripsy, *Urology Practice* (2016), doi: 10.1016/j.urpr.2016.04.002.

DISCLAIMER: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our subscribers we are providing this early version of the article. The paper will be copy edited and typeset, and proof will be reviewed before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to The Journal pertain.

All press releases and the articles they feature are under strict embargo until uncorrected proof of the article becomes available online. We will provide journalists and editors with full-text copies of the articles in question prior to the embargo date so that stories can be adequately researched and written. The standard embargo time is 12:01 AM ET on that date.

Running Head – Risk Factors for Readmission Following SWL

Risk Factors for Readmission Following Shock Wave Lithotripsy

Benjamin Nelson, MD; Anees Fazili, MD; Erdal Erturk, MD

University of Rochester Medical Center

Keywords: ESWL, Urolithiasis

Download English Version:

<https://daneshyari.com/en/article/8830358>

Download Persian Version:

<https://daneshyari.com/article/8830358>

[Daneshyari.com](https://daneshyari.com)