



Contents lists available at ScienceDirect

## The American Journal of Surgery

journal homepage: [www.americanjournalofsurgery.com](http://www.americanjournalofsurgery.com)

# Venous thromboembolism in common laparoscopic abdominal surgical operations

Reza Fazl Alizadeh, Sarath Sujatha-Bhaskar, Shiri Li, Michael J. Stamos, Ninh T. Nguyen\*

Department of Surgery, University of California, Irvine School of Medicine, USA

## ARTICLE INFO

### Article history:

Received 11 March 2017

Received in revised form

30 July 2017

Accepted 5 August 2017

### Keywords:

Venous thromboembolism

Laparoscopic abdominal operation

Pulmonary embolism

## ABSTRACT

**Background:** Venous thromboembolism (VTE) is potentially a serious postoperative complication. We examined the incidence and outcome of VTE among different laparoscopic abdominal surgical operations for benign diseases.

**Methods:** The National Surgical Quality Improvement Program database was utilized to evaluate all patients with benign disease who underwent laparoscopic abdominal operations including colorectal surgery, bariatric surgery, cholecystectomy, esophageal surgery, abdominal wall hernia repair, and appendectomy from 2005 to 2014. Multivariate logistic regression analysis was performed.

**Results:** 750,159 patients were studied and the overall incidence of VTE was 0.32% within 30 days of operation. Colorectal surgery had the highest incidence of VTE (734/65512, 1.12%) with significantly longest length of stay and operative time. Patients who developed VTE had higher mortality and worse outcomes compared to non-VTE patients.

**Conclusions:** Laparoscopic colorectal operations for benign disease is at higher risk for development of VTE compared to other laparoscopic abdominal operations. Further studies should be performed to elucidate the underlying mechanisms for our finding.

© 2017 Elsevier Inc. All rights reserved.

## 1. Introduction

Postoperative venous thromboembolism (VTE) is one of the most common post-surgical complications, which may have disastrous impacts on surgical outcomes.<sup>1</sup> VTE includes deep vein thrombosis (DVT) and pulmonary embolism (PE) and is one of the leading causes of morbidity and prolonged hospitalization in surgical patients,<sup>2,3</sup> and it has been accounted for 5–10% of all in-hospital deaths in US.<sup>4,5</sup>

Although, several VTE preventative guidelines and recommendation have been issued from different surgical societies,<sup>6,7</sup> recent publications demonstrated that there is an incremental trend for VTE occurrences.<sup>8</sup> This increase of incidences is most likely related to the proliferation of higher risk patients who undergo surgical procedures, varied outcomes, and low rates of compliance with VTE

preventative guidelines.<sup>9,10</sup>

Since the introduction of laparoscopy, minimally invasive techniques have been adopted across multiple surgical disciplines and is now the preferred surgical approach of choice for most common abdominal operations.<sup>11</sup> Certain factors associated with laparoscopy may contribute to a higher risk for VTE, such as the use of pneumoperitoneum, reverse Trendelenburg positioning, and prolonged operative times; however, recent published studies revealed that laparoscopic surgery is associated with a lower incidence of VTE compared with that of open surgery in several commonly performed gastrointestinal procedures.<sup>12,13</sup> The aim of this study was to determine the incidence of VTE rates among different laparoscopic abdominal surgical procedures. Since malignancy has been well shown to be an independent risk factor for VTE,<sup>14,15</sup> we select our patient population that underwent abdominal surgery specifically for benign diseases to avoid malignancy being a confounding variable.

## 2. Materials and methods

The American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) database is the first nationally

*Abbreviations:* VTE, Venous thromboembolism; DVT, Deep vein thrombosis; PE, Pulmonary embolism; NSQIP, National Surgical Quality Improvement Program; SSI, Superficial surgical site infection; AOR, Adjusted odds ratio; CI, Confidence interval.

\* Corresponding author. Department of Surgery, University of California, Irvine School of Medicine, 333 City Blvd West, Suite 1600, Orange, CA, 92868, USA.

E-mail address: [ninhn@uci.edu](mailto:ninhn@uci.edu) (N.T. Nguyen).

<http://dx.doi.org/10.1016/j.amjsurg.2017.08.032>

0002-9610/© 2017 Elsevier Inc. All rights reserved.

validated, outcomes-based program designed for the purpose of improving surgical quality of care. It is comprised of more than 500 hospitals that collect hospital-level data on preoperative risk factors, intraoperative variables, and 30-day morbidity and mortality after inpatient and outpatient surgical procedures. A trained surgical clinical reviewer is used at participating hospitals to standardize data capture and ensure reliability.<sup>16</sup>

The 2005 to 2014 ACS-NSQIP database was retrospectively reviewed to identify patients who underwent laparoscopic abdominal operations. Approval for the use of the NSQIP patient-level data in this study was obtained from the institutional review board of the University of California, Irvine Medical Center and NSQIP.

We examined all patients who underwent laparoscopic abdominal operations based on the following Current Procedural Terminology (CPT) codes: cholecystectomy (47562, 47563, 47564), esophageal surgery (gastroesophageal reflux: 43279, 43280, 43283, paraesophageal hernia: 43281, 43282), bariatric surgery (43644, 43645, 43659, 43770, 43771, 43772, 43773, 43774, 43775), colorectal surgery (colectomy: 44204, 44205, 44206, 44207, 44208, 44210, 44211, 44212, proctectomy: 45395, 45397, 45400, 45402, 45499), hernia repair (inguinal hernia: 49650, 49651, incisional hernia: 49652, 49653, 49654, 49655, 49656, 49657), and appendectomy (44970, 44979). Patients who had malignant diseases and patients younger than 18 years were excluded from this study.

All the patients with VTE were identified and divided into six groups based on sub-specialty: colorectal surgery, bariatric surgery, cholecystectomy, esophageal surgery, hernia repair, and appendectomy. In our subset analysis, patients were divided into VTE vs. non-VTE groups. Postoperative VTE was defined according to NSQIP original variables of deep vein thrombosis (DVT) and pulmonary embolism (PE). Variables used in the analysis were provided by the NSQIP database and included demographic data (age, race, and gender), comorbidities, operation length, type of procedures, type of admission (emergent vs. non-emergent), length of stay, and postoperative complications.

The outcomes evaluated were 30-day mortality, superficial surgical site infection (SSI), organ space SSI, deep incisional SSI, wound disruption, pneumonia, unplanned intubation, ventilator dependency more than 48 h, urinary tract infection, acute renal failure, renal insufficiency, postoperative sepsis, postoperative septic shock, and bleeding disorders. Risk adjusted analysis was performed to identify predictors of VTE.

### 2.1. Statistical analysis

Statistical analyses were conducted using SPSS software, Version 23 (SPSS Inc., Chicago, IL). The main analysis method was multivariate analysis using logistic regression model to explore the independent association between perioperative factors and complications of VTE. In order to report risk adjusted data, adjustments were made for all the variables of the study. For each variable, the adjusted odds ratio (AOR) with a 95% confidence interval was calculated. Statistical significance was set at *P* values greater than 0.05.

### 3. Results

A total of 750,159 laparoscopic abdominal operations were performed with 2424 (0.32%) patients diagnosed with VTE within 30 days of operation. Of these, the VTE rate was highest for colorectal surgery at 1.12% (734 of 65,512) and lowest following appendectomy at 0.16% (276 of 168,963). The VTE rate was 0.5% following esophageal surgery (171 of 31,511), 0.35% following bariatric surgery (539 of 153,552), 0.2% following cholecystectomy

(489 of 239,499), and 0.24% following incisional and inguinal hernia repair (215 of 91,122) (Fig. 1). The median age of patients with VTE was 59 years for colorectal surgery; 48 years for bariatric surgery; 63 years for cholecystectomy; 68 years for esophageal surgery; 61 years for incisional and inguinal hernia repair; and 51 years for appendectomy.

Patient demographics and comorbidities for patients with VTE according to different laparoscopic abdominal operation are listed in Table 1. For patients with VTE, mean length of stay ( $13 \pm 11.5$  days) and operative time ( $209 \pm 111$  min) were longer for colorectal operations compared to other abdominal operations (Table 1).

In the subset analysis, patients with VTE and non-VTE were studied. Patient demographics and comorbidities of patients with VTE vs. non-VTE are listed in Table 2. Mean length of hospitalization was significantly longer in patients with vs. without VTE ( $8.4 \pm 11$  vs.  $2 \pm 6$  days, respectively,  $P < 0.0001$ ). Mean operation time in patients with and without VTE were  $140 \pm 99$  and  $85 \pm 62$ , respectively ( $P < 0.0001$ ) (Table 2).

Risk adjusted analysis of top four factors associated with VTE includes chronic steroid use (AOR 2.71,  $P < 0.01$ ), preoperative blood transfusion (AOR 2.70,  $P < 0.01$ ), Black or African American ethnicity (AOR 2.69,  $P < 0.01$ ), and preoperative sepsis (AOR 2.55,  $P < 0.01$ ) (Table 3).

The unadjusted mortality rates in patients with and without VTE were 3.3 and 0.2%, respectively. After adjusting for comorbidities, risk of mortality was higher in patients with VTE (AOR 4.54, CI 3.40–6.05,  $P < 0.0001$ ). Patients with VTE following colorectal had the highest risk for mortality compared to other procedures (AOR 2.36, CI 1.89–2.94,  $P < 0.0001$ ). Multivariate analysis also confirmed the association of VTE with worse patient outcomes. Ventilator dependency (AOR 10.51, CI 8.22–12.52,  $P < 0.0001$ ), unplanned re-intubation (AOR 9.76, CI 8.07–11.80,  $P < 0.0001$ ), renal insufficiency (AOR 8.17, CI 6.0–11.13,  $P < 0.0001$ ), pneumonia (AOR 8.22, CI 6.86–9.84,  $P < 0.0001$ ), and septic shock (AOR 7.46, CI 5.90–9.42,  $P < 0.0001$ ) were the most significantly increased post-surgical outcomes (Table 4).

Finally, on multivariate analysis, a comparison of the likelihood of developing VTE following several laparoscopic abdominal procedures demonstrated the odds of developing VTE following cholecystectomy (AOR: 0.94; 95% CI: 0.75–1.17;  $p = 0.60$ ) and hernia repair (AOR: 1.0; 95% CI: 0.80–1.26;  $p = 0.95$ ) were not significantly different from the baseline of appendectomy. However, odds were significantly higher in patients undergoing colorectal surgery (4.99; 3.99–6.24;  $p < 0.0001$ ), bariatric surgery (1.36; 1.01–1.82;  $p < 0.037$ ), and esophageal surgery (1.80; 1.31–2.47;  $p < 0.0001$ ) (Fig. 2).

### 4. Discussion

There are a variety of risk factors for development of VTE in surgical patients. Some of these factors include the type of surgical procedure, the degree of invasiveness, type of anesthesia, and extended period of immobilization.<sup>17,18</sup> With significant growth of minimally invasive surgery in last few decades, laparoscopic surgery is the procedure of choice for most abdominal procedures. The risk for VTE complications after laparoscopic surgery has been considered potentially higher than that of open surgery because of the use of pneumoperitoneum, altered patient positioning and possibly a prolonged operative time during the learning curve of the laparoscopic procedure.<sup>19–21</sup> However, recent studies have demonstrated that the overall incidence of VTE is actually lower after laparoscopic operation compared with open operation.<sup>12,13,22</sup> In the current study, we examined the incidence of VTE among six laparoscopic abdominal surgical procedures and found that colorectal surgery have the highest incidence of postoperative VTE. The

Download English Version:

<https://daneshyari.com/en/article/8830913>

Download Persian Version:

<https://daneshyari.com/article/8830913>

[Daneshyari.com](https://daneshyari.com)