

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.e-asianjournalsurgery.com

TECHNICAL NOTE

An unusual salvage technique for posterior tracheal membranous laceration associated with transhiatal esophagectomy: A transcervical–transsternal approach

Seyed Ziaeddin Rasihashemi ^a, Nassir Rostambeigi ^b,
Seyed Reza Saghebi ^c, Faeze Daghigh ^d, Ali Ramouz ^{e,*}

^a Department of Cardiothoracic Surgery, Tabriz University of Medical Sciences, Tabriz, Iran

^b Department of Radiology, University of Minnesota Medical Center, Minneapolis, MN, USA

^c Tracheal Diseases Research Center, National Research Institute of Tuberculosis and Lung Diseases (NRITLD), Shahid Beheshti University of Medical Sciences, Tehran, Iran

^d Student Research Committee & Medicine Faculty, Daneshgah street, 5183915881, Tabriz University of Medical Sciences, Tabriz, Iran

^e Student Research Committee & Medicine Faculty, Tabriz University of Medical Sciences, Tabriz, Iran

Received 9 November 2015; received in revised form 23 December 2015; accepted 24 December 2015

KEYWORDS

esophagectomy;
laceration;
trachea;
transcervical;
transhiatal

Summary Various surgical approaches may be employed for esophageal resection. Major airway injuries due to transhiatal esophagectomy include vertical tears in the membranous trachea. Tracheal injury is an uncommon but potentially fatal complication. This article describes the technique to repair the posterior membranous tracheal tear, extended just over the carina through a transcervical–transsternal approach, thereby avoiding a second thoracotomy. Six patients with posterior membranous tracheal injury underwent this procedure. The laceration ranged from 3 cm to 5 cm in length. Four patients had received neoadjuvant chemoradiation. The management of tracheal laceration added approximately 60 minutes to the total operation time. There was no mortality related to tracheal injury. Patients were followed up for 6 months after surgery, and both posterior tracheal wall and transverse tracheotomy remained intact without stenosis. The transcervical–transsternal approach decreases the need of thoracotomy and its complications in patients with tracheal laceration in any stage, even in cases of an extended tear down to the carina.

Conflicts of interest: The authors declare no conflicts of interest.

* Corresponding author. Student Research Committee & Medicine Faculty, Daneshgah street, 5183915881, Tabriz University of Medical Sciences, Tabriz, Iran.

E-mail address: ali.ramouz@gmail.com (A. Ramouz).

<http://dx.doi.org/10.1016/j.asjsur.2015.12.009>

1015-9584/Copyright © 2016, Asian Surgical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Rasihashemi SZ, et al., An unusual salvage technique for posterior tracheal membranous laceration associated with transhiatal esophagectomy: A transcervical–transsternal approach, Asian Journal of Surgery (2016), <http://dx.doi.org/10.1016/j.asjsur.2015.12.009>

Copyright © 2016, Asian Surgical Association. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Transhiatal esophagectomy (THE) is a widely used approach with low rates of morbidity and mortality. However, the proximity of the esophagus to vital organs within the chest increases the possibility of inadvertent injury during blunt dissection.^{1,2} Membranous tracheal and bronchial injuries are potentially lethal complications of blunt transhiatal mobilization of the esophagus due to the close vicinity of the esophagus to these structures.³ With due attention to the prevalence of THE among surgeons, the incidence of tracheal injuries has been reduced to less than 10%, and even to less than 2% in some reports from experienced centers.^{3,4}

Major airway injuries due to THE are usually vertical tears in the posterior membrane of the trachea. Various strategies in the management of tracheal lacerations have been reported. Although thoracotomy is the main part in many of these procedures, a variety of technical modifications have been reported to repair tracheal laceration in THE procedures without thoracotomy. We aimed to describe a transcervical–transsternal method for repair of membranous tracheal lacerations following THE.^{2–5}

2. Technique

A retrospective study was performed on all patients who underwent THE in our general tertiary hospital over a period of 11 years (2000–2012). Of the 782 patients, six sustained an iatrogenic injury in the posterior membrane of the trachea. All tears were recognized intraoperatively. The study group included four women and two men, with a mean age of 59 years (range, 52–70 years).

In five cases, the squamous cell carcinoma was located in the middle third of the esophagus, and one patient had carcinoma distal to the carina. All cases had complaints of dysphagia and retrosternal pain, lasting between 3 months and 8 months. The TNM staging was II in four patients and III in two patients. Four patients had a history of neoadjuvant chemoradiation prior to surgery (Table 1).

Bronchoscopy was performed in patients with cancer of the middle third of the esophagus; the posterior wall of the tracheas was not implicated/infiltrated by the tumor.

During cervical dissection, tears were recognized in the posterior membranous of the trachea near the thoracic inlet. The laceration ranged from 3 cm to 5 cm in length. Tracheal rupture was manifested immediately with a fall in airway resistance and ventilator air leakage through the incision site. In order to prevent progressive pressure drop, the tube cuff deflated immediately, and the tube was advanced to the level of the carina. Despite the tracheal tube advancement, O₂ saturation dropped. Therefore, we performed an anterior transverse tracheotomy between

two cartilaginous rings above the laceration site (Figure 1). The orotracheal tube was withdrawn, and a 6-mm (ID) sterile, low-pressure cuffed, flexible armored endotracheal tube was inserted distal to laceration through the tracheal incision. O₂ saturation and respiratory rate returned to normal ranges. Owing to inadequate exposure to the whole mediastinal trachea through the classic anterior sternocleidomastoid incision, we extended the incision and performed a partial sternal split to the third intercostal space with manubrial retraction. All patients underwent esophagectomy prior to repair of the tracheal rupture. The lateral tracheal blood supply and both recurrent laryngeal nerves were preserved, and the laceration was repaired from distal to proximal with absorbable interrupted sutures outside of the tracheal lumen. While completing the repair of posterior laceration, the endotracheal tube was removed intermittently, and subsequently, the transverse

Table 1 Demographic data with diagnostic features and intraoperative characteristics of the patients.

	Frequency (%)	Mean ± SD	Range
Age (y)		59 ± 7.8	50–70
Sex			
Male	2 (33)		
Female	4 (66)		
Pathology			
SCC	6 (100)		
Adenocarcinoma	0		
TNM staging			
I	0		
II	4 (66)		
III	2 (33)		
IV	0		
Tumor location			
Upper 1/3	0		
Middle 1/3	5 (83)		
Lower 1/3	1 (17)		
Operative time (min)		198 ± 25	165–240
Distance to tumor proximal end (cm)		11 ± 2	8–16
Tracheal injury length (mm)		25 ± 5	17–34
Rupture repair duration (min)		49 ± 5	42–58
Treatment			
Surgery	3 (50)		
Chemoradiotherapy	0		
Surgery and chemoradiotherapy	3 (50)		

SCC = squamous cell carcinoma; SD = standard deviation.

Download English Version:

<https://daneshyari.com/en/article/8831064>

Download Persian Version:

<https://daneshyari.com/article/8831064>

[Daneshyari.com](https://daneshyari.com)